

PH 15 572-246 1/2  
1-29-90

PANTONE®  
179C

# Rockland. C-EM-DIE

TRADE MARK REGISTERED

DAIRY, LIVESTOCK & BARN

## CONTACT & SPACE SPRAY

WITH QUICK KNOCKDOWN AND KILL CHEMICALS

KILLS Stable Flies-Horn Flies-House Flies-Mosquitoes and Chicken Mites-When used as directed



<b>ACTIVE INGREDIENTS:</b>	
*Tech. Piperonyl Butoxide .....	0.277%
2,2-dichlorovinyl dimethyl phosphate .....	0.230%
Related Compounds .....	0.020%
Pyrethrins .....	0.034%
**INERT INGREDIENTS .....	99.439%
	100.000%

\*Equivalent to .22% (butylcarbityl)(6-propylpiperonyl) ether  
and .55% related compounds  
\*\*Contains Petroleum Distillate

EPA Reg. No. 572-246


EPA Est. No. 572-NJ-1

**KEEP OUT OF REACH OF CHILDREN**

**CAUTION:**

SEE SIDE PANEL FOR ADDITIONAL CAUTIONS  
NET CONTENTS 1 GALLON

Not Reviewed. Registrant claims to be in accordance with  
Reregistration of Pesticide Product Guidance for Phase 2  
Response, Page 2.12, Active to Inert Change in Status.



C-EM-DIE Dairy and Barn Spray is a special formulation compounded primarily for the dairy industry. This product will not destroy the metal of any sprayer; no waste or sediment in can.

7/2

### PRECAUTIONARY STATEMENTS

**HAZARDS TO HUMANS AND DOMESTIC ANIMALS:** Harmful if swallowed, inhaled or absorbed through the skin. Wash thoroughly with soap and water after handling and before eating or smoking. Do not contaminate utensils, feed, water or foodstuffs. Avoid breathing of spray mist. Avoid wetting the hides of animals.

Do not use this product on animals simultaneously or within a few days before or after treatment with or exposure to cholinesterase inhibiting drugs, pesticides, or chemicals. Do not apply in combination with trichlorofon. Do not use on Brahman or Brahman cross cattle, as they may show hypersensitivity to organic phosphates.

**ENVIRONMENTAL HAZARDS:** This product is toxic to fish, birds and other wildlife. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

**PHYSICAL OR CHEMICAL HAZARDS:** DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME.

### DIRECTIONS FOR USE FOLLOW THOROUGHLY

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Spray, at the rate of 1-2 oz. per 100 sq. ft., on walls, ceilings, pens, posts, stalls, windows, doors, where flies congregate; holding sprayer about one foot from surface being sprayed and close doors and windows for 10 minutes when spraying to avoid draft and waste.

**CATTLE, CALVES, HOGS, PIGS:** Spray animals with a fine mist once a day to kill horn flies, houseflies, mosquitoes. **DAIRY CATTLE-**for killing these flies, spray lightly 20-30 minutes before or after milking time and only when necessary. No more than 2 ounces of spray should be applied per animal per application. For young animals, apply no more than one ounce per animal as a spray. Tobeinids to be killed must be hit directly by spray.

Wash the teats of sprayed dairy animals with an approved cleansing solution and then dry them prior to milking.

**CHICKEN MITES:** Empty nests before spraying. Spray roosts, at the rate of 1-2 oz. per 100 sq. ft., paying particular attention to all cracks and crevices and other hidden places. Thoroughly clean the premises before spraying as this will help in controlling mites. Repeat application in about 2 weeks if necessary. Spray entire interior of the coop and house thoroughly.

### BARN SPRAY:

Use C-EM-DIE in barns, milk houses, outbuildings, pens for effective control of Stable Flies, House Flies, Horn Flies. Use powder or compressed air sprayer, if preferred, that throws a good fine even spray. Animals should be removed from barn before treatment. Spraying of animals should be done outdoors as a separate treatment. Direct the mist toward the upper portion of the enclosure, filling the air with fine mist, using 2 ounces of mist per 1000 cubic feet of air space. For best results keep doors and windows closed when possible for 10 minutes after application and during spraying. Applicator must not remain in treated area. After the 10 minute treatment period, the area should then be opened up and ventilated prior to reoccupying by humans or animals. Do not contaminate feed, drinking water, milk or milking equipment. Do not apply in areas where animals have received direct application of DDVP within 8 hours.

### STORAGE AND DISPOSAL


**STORAGE:** Store unused material, in original container only, in a locked, dry storage area out of reach of children and animals. Do not contaminate water, food or feed by storage or disposal. Do not store near heat or open flame.

**PESTICIDE DISPOSAL:** Pesticide wastes are toxic. Improper disposal of excess pesticides, spray mixture or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest E.P.A. Regional Office for guidance.

**CONTAINER DISPOSAL:** Do not reuse empty container. Securely wrap closed, original container in several layers of newspaper and discard in trash.

**WARRANTY:** Seller makes no warranty, expressed or implied concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

Manufactured by

 Rockland Chemical Co., Inc.  
West Caldwell, NJ 07006 U.S.A.

Product C24601  
E.P.A. Est. 572-NJ-1  
EPA Reg. No. 572-246