

ACCEPTED  
12-2-71  
572-248

**LETTUCE:** Downy mildew — in plant bed, use as drenching spray starting when seed is planted and continue at 7 to 10 day intervals until transplanting. In field, begin applications when disease appears and continue at 7 to 10 day intervals as long as needed (under severe conditions, shorten interval to 3 to 5 days). Do not apply later than 7 days before harvest.

**COMPATIBILITY:** ROCKLAND ZINEB GARDEN FUNGICIDE is compatible with common insecticides, such as methoxychlor, organophosphorus compounds, nicotine sprays, arsenicals and rotenone.

**CAUTION:** May cause irritation of eyes, nose, throat, and skin. May be harmful if inhaled or swallowed.

Avoid breathing dust or spray mist.

Avoid contact with skin, eyes and clothing.

Keep away from fire or sparks.

In case of contact, flush with plenty of water; for eyes, get medical attention.

Do not reuse container. Destroy when empty.

When used as directed on label and in accordance with good practices, this material complies with the Food and Drug Law and other Federal Laws.

**WARRANTY:** Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

EPA Reg. No. 572-248



ROCKLAND

NET WT. 8 OUNCES

**ROCKLAND  
ZINEB  
GARDEN FUNGICIDE**

**ACTIVE INGREDIENT:**

Zineb (Zinc ethylenebisdithiocarbamate) 75.0%  
(Total Zinc as Metallic 17.7%)

**INERT INGREDIENTS:** ..... 25.0%

**DIRECTIONS FOR USE**

**FOR FLOWERS,** use sufficient spray to provide thorough coverage. Be sure to spray underside of leaves. Mix 2 ounces ROCKLAND ZINEB GARDEN FUNGICIDE in 8 gallons of water or 1½ table-spoons per gallon. Mix well so Zineb is dissolved thoroughly. Spray every 7 to 10 days to control: (Rust and Stem Rot on Asters; Leaf Spot on Carnations, Iris, Pansy, Zinnia; Botrytis Blight on Dahlia, Hydrangea, Peony; Gray Mold on Geranium, Lily; Rust, Powdery Mildew, Botrytis Blight on Snapdragon; Leaf Spot on Gladiolus) and Botrytis Blight on Tulips.)

**FOR VEGETABLE CROPS,** mix 2 ounces in 8 gallons of water or 8 ounces in 32 gallons of water. Spray thoroughly.

**TOMATOES:** Late and early blights, Septoria leaf spot, gray leaf spot (Stemphylium), anthracnose — start applications when fruit of first clusters is well formed; repeat at 7 to 10 day intervals. When weather conditions favor late blight, more frequent applications may be necessary, starting when plants are set in the fields.


Apply to dewy or wet plant bed using  
 spraying spray starting when seed is planted and  
 repeat at 7 to 10 day intervals until transplanting  
 field near hot spots when disease appears  
 and control at 7 to 10 day intervals as long as  
 plants and weather conditions shorten interval  
 to 5 days; Do not apply later than 7 days  
 before harvest.

**COMPATIBILITY.** ROCKLAND ZINEB GARDEN  
 FUNGICIDE is compatible with common insecti-  
 cides such as malathion, DDT, organophosphorus com-  
 pounds, and many other insecticides and fungicides.

**CAUTION.** May cause irritation of eyes, nose,  
 throat, and skin. Avoid contact with skin and or  
 clothing.



**ROCKLAND**



- ① GARDEN FUNGICIDE.
- ② FOR CERTAIN PLANT DISEASES.
- ③ ON FLOWERS AND VEGETABLES.

**CAUTION: KEEP OUT OF REACH OF CHILDREN**  
**See back panel for additional cautions**

**ROCKLAND CHEMICAL CO., INC.**  
 Passaic Avenue, West Caldwell, N. J.

ROCKLAND CHEMICAL CO., INC.  
 1000 WEST CALDWELL AVENUE  
 WEST CALDWELL, N. J.

**DIRECTIONS FOR USE**

Apply to dewy or wet plant bed using  
 spraying spray starting when seed is planted and  
 repeat at 7 to 10 day intervals until transplanting  
 field near hot spots when disease appears  
 and control at 7 to 10 day intervals as long as  
 plants and weather conditions shorten interval  
 to 5 days; Do not apply later than 7 days  
 before harvest.