


ROCKLAND CHEMICAL CO., INC.
P. O. BOX 809
WEST CALDWELL, N. J. 07007-0809

FRONT PANEL:



Rockland
Zineb
Garden
Fungicide

Controls diseases on vegetables and ornamentals
Compatible with common insecticides
Use 1½ Tablespoons per 1 gallon of water

ACTIVE INGREDIENT:


Zineb (Zinc ethylenebisdithiocarbamate)..... 75.0%
(Total Zinc as Metallic..17.7%)

INERT INGREDIENTS:..... 25.0%
100.0%

14 point type

CAUTION: KEEP OUT OF REACH OF CHILDREN *← 14 point type*
See back panel for additional cautions.

Net Weight EIGHT OUNCES



ROCKLAND CHEMICAL CO., INC.

P. O. BOX 809

WEST CALDWELL, N. J. 07097-0809

BACK PANEL:

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS:

CAUTION: MAY CAUSE IRRITATION OF EYES, NOSE, THROAT AND SKIN. MAY BE HARMFUL IF INHALED OR SWALLOWED. Do not breathe dust or spray mist. Protective clothing consisting of a long sleeved shirt, gloves, hat and boots must be worn during mixing and applying.

Do not feed forage to dairy animals or animals being finished for slaughter.

STATEMENT OF PRACTICAL TREATMENT:

IF ON SKIN: Wash with water.

IF IN EYES: Wash copiously with water and get medical attention.

ENVIRONMENTAL HAZARDS:

This pesticide is toxic to aquatic organisms. Do not apply directly to water. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by the cleaning of equipment or disposal of wastes. PHYSICAL HAZARDS: Keep away from fire and sparks.

This product is highly toxic to bees exposed to direct treatment or residues on plants. Protective information may be obtained from your Cooperative Agricultural Extension Service.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Use only in accordance with recommendations on this label.

COMPATIBILITY: ROCKLAND ZINEB GARDEN FUNGICIDE is compatible with common insecticides, such as Methoxychlor, organophosphorus compounds, nicotine sprays, arsenicals and Rotenone.

The addition of ROCKLAND SPREADER-STICKER to the spray mixture improves distribution of the spray.

Do not contaminate water, food or feeds by storage or disposal.

STORAGE: Store in cool, dry place. Do not allow to become wet or overheated during storage. This may bring on chemical changes which will impair its effectiveness and may also generate flammable vapors. Keep container closed when not in use. DISPOSAL: Do not re-use empty container. Wrap container and put in trash collection.

FOR FLOWERS: Use sufficient spray to provide thorough coverage. Be sure to spray underside of leaves. Mix 2 ounces in 8 gallons of water (1½ Tablespoons per gallon). Mix well so Zineb is dissolved thoroughly. Spray every 7 to 10 days to control: Rust and Stem Rot on Asters; Leaf Spot on Carnations, Iris, Pansy, Zinnia; Botrytis Blight on Dahlia, Hydrangea, Peony; Cladosporium Gray Mold on Geranium, Lily; Rust, Powdery Mildew, Botrytis Blight on Snapdragon; Leaf Spot on Gladiolus and Botrytis Blight on Tulips.

ROCKLAND CHEMICAL CO., INC.

P. O. BOX 229

WEST CALDWELL, N. J. 07007-0229

FOR VEGETABLE CROPS: Mix 2 ounces in 8 gallons of water (8 ounces in 32 gallons). Spray thoroughly.

TOMATOES: Anthracnose, Cladosporium Gray Mold, Early and Late Blight, Septoria Leaf Spots and Stemphylium. Apply when disease appears and repeat at 7 to 10 day intervals or more often if disease is severe. Do not apply within 5 days of harvest.

BEANS: Anthracnose, Downy Mildew, Rust. Begin when first blossoms appear and repeat at 7 to 14 day intervals. Do not apply within 7 days of harvest.

CORN (Sweet): Helminthosporium Leaf Blight, Puccinia Rust. Apply when disease appears and repeat at 3 to 7 day intervals. Do not feed treated forage to dairy animals or animals being finished for slaughter.

LETTUCE: Downy Mildew - in plant bed, use as drenching spray starting when seed is planted and continue at 7 to 10 day intervals until transplanting.

In field, begin applications when disease appears and continue at 7 to 10 day intervals as long as needed (under severe conditions, shorten interval to 3 to 5 days). Do not apply later than 10 days before harvest.

Tablespoon enclosed for easy measuring.

This package is sold by weight, not by volume. Some settling of contents may have occurred during shipping and handling.

WARRANTY: Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

Manufactured by

Rockland Chemical Co., Inc.
West Caldwell, NJ 07006 U.S.A.

Product

EPA Reg. No. 572-248

EPA Est. 572-NJ-1

