

**CAUTION: MAY IRRITATE EYES, NOSE, THROAT AND SKIN.**

Avoid breathing dust. Avoid contact with skin, eyes, and clothing. Wash thoroughly after using. Keep away from fire or sparks. In case of contact flush skin or eyes with plenty of water; for eyes, get medical attention.

**IMPORTANT** — Never allow ROCKLAND BENOMYL FUNGICIDE to become wet during storage. This may lead to certain chemical changes which will reduce the effectiveness of ROCKLAND BENOMYL FUNGICIDE as a fungicide. Keep container closed when not in use. Do not re-use container. Bury when empty.

**DIRECTIONS**

ROCKLAND BENOMYL FUNGICIDE is recommended for the control of certain diseases of roses, flowers, ornamentals, and turf. It is highly effective for the control of many fungus diseases, providing both curative (eradicator) action and protective action. ROCKLAND BENOMYL FUNGICIDE is a wettable powder to be mixed with water for application as a spray. Addition of a surfactant to the spray mixture enhances curative action of the fungicide and improves distribution of the spray on hard-to-wet plants such as roses.

**ROSES** (Field and Greenhouse) — **Powdery Mildew, Black Spot:** Use one tablespoonful per 2 gals. water. Begin application when disease first appears and repeat at 10 to 14 day intervals throughout the growing season. Shorten intervals during humid rainy weather.

**FLOWERS, ORNAMENTALS** (Field and Greenhouse) — **Powdery Mildew, Botrytis Gray Mold:** Use one tablespoon per 2 gals. water. Begin application when disease first appears and repeat at 10 to 14 day intervals as long as needed.

**TURF** — **Dollar Spot** (*Sclerotinia homeocarpa*), **Large Brown Patch** (*Rhizoctonia solani*): Apply at the rate of 2 oz. (12 tablespoonfuls) per 1000 sq. ft. in sufficient water to obtain thorough coverage, usually 2 to 5 gals. per 1000 sq. ft. Apply at earliest appearance of disease and continue at 10 to 14 day intervals as long as needed. When conditions are unusually favorable for development of disease, reduce interval to 5 to 7 days. Do not graze or feed clippings from treated areas to livestock.

**STRIPE SMUT** — Make a single application of ROCKLAND BENOMYL FUNGICIDE at the rate of 6 oz. per 1000 sq. ft. (3 oz. per 500 sq. ft.) in October or in early spring before grass begins growth period. Use sufficient water to obtain thorough coverage, usually 2 to 5 gallons per 1000 sq. ft. (1 to 3 gals. per 500 sq. ft.) After application, water turf sufficiently to carry product to base of plants.

**WARRANTY:** Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

Packaged and Distributed by  
**ROCKLAND CHEMICAL CO., INC.**  
PASSAIC AVENUE WEST CALDWELL, N. J.




Net Contents: 4 oz.

**ROCKLAND  
BENOMYL  
FUNGICIDE**

WETTABLE POWDER

For Turf,  
Flowers, Roses,  
Ornamentals

ACCEPTED  
3-3-72  
572-259  
GENERAL INVESTIGATIONS  
FOR PATENTED CLAIMS



**CAUTION: KEEP OUT OF REACH OF CHILDREN.**

ACTIVE INGREDIENT:

\*Dexamyl (Methyl 1-(butylcarbamoyl)-2-benzimidazolecarbamate) ..... 50%

INERT INGREDIENTS: ..... 50%

\*U. S. Pat. Pending E. I. duPont de Nemours and Company EPA Reg. No. 572-254

SEE BACK PANEL FOR ADDITIONAL CAUTIONS.