

PM 25

572-255

4/28/93

P5172

ACCEPTED

APR 28 1993


Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under

572-255

Feed & Seed

18-6-6 with

Tupersan®



Treats 5,000 Sq. Ft.
Newly Seeded Lawn

Apply After Seeding
Controls Crabgrass
& Fertilizes



**Feed & Seed 18-6-6
with Tupersan®**

| | |
|--|---------|
| ACTIVE INGREDIENT | |
| Souron [1-(2-methylcyclohexyl)-3-phenylurea] | 1.50% |
| INERT INGREDIENTS | 98.50% |
| | 100.00% |

® Registered Trademark of E. I. duPont de Nemours & Co. for Souron Weed Killer.

EPA Reg. No. 572-255 EPA EST. No. 9198-01-1

KEEP OUT OF REACH OF CHILDREN
CAUTION

See back panel for complete Precautionary Statements.

| | |
|--|-----|
| GUARANTEED ANALYSIS | |
| Total Nitrogen (N) | 18% |
| 15.6% Urea Nitrogen | |
| 2.4% Ammoniacal Nitrogen | |
| Available Phosphoric Acid (P ₂ O ₅) | 6% |
| Water Soluble Potash (K ₂ O) | 6% |

Primary Nutrient Sources: Ammoniated Phosphorus, Urea, Murate of Potash

Potential Acidity equivalent to 67.5 lbs. Calcium Carbonate per ton.

NET WT 40 LBS (70 L)

272

ROCKLAND FEED AND SEED is a very effective lawn fertilizer and crabgrass control compound. These materials normally are applied during the spring of the year and, by combining them in this manner, treatments are applied in one application with a direct saving in time and money.

The Tupersan in this product is highly effective for the control of crabgrass (smooth and hairy) and certain annual weed grasses in turf areas such as golf course fairways, lawns, parks, roadsides and turf grown for grass seed or sod production. Because of its high degree of selectivity, ROCKLAND FEED AND SEED may be applied at time of seeding without causing injury to germinating seeds of many common temperate region grasses as well as established turf.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS: Harmful if swallowed. Avoid contact with eyes, skin or clothing. Avoid breathing dust. In case of eye contact, flush eyes with plenty of water. Wash thoroughly after using. Keep children and pets off treated areas until this material is washed into the soil and grass is dry. Do not contaminate feed or foodstuffs.

ENVIRONMENTAL HAZARDS: This product is toxic to fish. Do not apply directly to water to areas where surface water is present or to Intertidal areas below the high water mark. Do not contaminate water when disposing of equipment washwaters.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. USE ONLY ON TURF AREAS. DO NOT apply to crops used for food or feed purposes.

ROCKLAND FEED AND SEED is recommended for pre-emergence control of annual weed grasses, such as crabgrass (smooth and hairy), foxtail, and barnyardgrass, in newly seeded or established plantings of the following grasses only: bluegrass, fescue, redtop, smooth brome, perennial ryegrass, orchardgrass, and the following strains of bentgrass: Penncross, Seaside, Highland Astoria, Nimisila, C-1, C-7, and C-19. Do not use on other bentgrass strains nor on Bermudagrass, as injury to turf may result. Treatment will not control annual bluegrass (*Poa annua*), clover or most broadleaf weeds. This product may be applied at any stage of turf development, including new seedings.

Before making application lawn should be raked to remove extraneous material and insure uniform application. Use any spreader in good condition. Measure off area to be treated. Do not spread over larger area than recommended. Water must be provided by rainfall or normal lawn watering within 3 days after application.

Established Turf and New Fall Seedings: Apply contents of this bag over 2500 square feet of lawn area in the spring.

New Spring Seedings: Apply contents of this bag over 5000 square feet of lawn area as final operation following seeding.

SUGGESTED SPREADER SETTINGS

| SPREADER MODEL | ESTABLISHED and FALL SEEDED LAWNS | NEWLY SEEDED LAWNS 5000 Sq. Ft. |
|----------------|-----------------------------------|---------------------------------|
| SCOTT | 7 1/2 | 4 1/2 |
| JACKSON | 7 1/2 | 5 1/2 |
| MODERN | 8 1/2 | 5 1/2 |
| CYCLONE | 4.3 | 3 |

These spreader settings are meant as a guide. They are in no sense a guarantee of exact application, inasmuch as age, wear of spreader, speed of operation and roughness of terrain can cause variations in rates applied.

STORAGE AND DISPOSAL

STORAGE: Do not store near seed, fertilizers and other pesticides. Store unused material in original container only, in a locked, dry storage area out of reach of children and animals. Do not contaminate water, food or feed by storage or disposal.

DISPOSAL: Materials that cannot be used according to label instructions, must be disposed of according to applicable Federal, State or local procedures. Do not reuse empty bag. Securely wrap in several layers of newspaper and discard in trash.

WARRANTY: Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

EPA REG. No. 572-255
#A25516

EPA Est. No. 9198-QH-1

MANUFACTURED FOR

