

FOR USE BY LAWN DOCTOR FRANCHISEES ONLY

A Balan-coated corn cob for pre-emergence control of crabgrass and annual blue grass in established lawns.

DIRECTION FOR USE

LAWN and ORNAMENTAL TURFS (established turf only): Apply at rate of 2 lbs. per 1,000 sq. ft. of lawn with a spreader for the control of Smooth and Hairy Crabgrass, Goosegrass (Silver Crabgrass), Barnyardgrass, Yellow and Green Foxtail and Annual Bluegrass (Poa Annua). This product controls weeds by killing their seeds as they germinate. It does not control established weeds. Time application in the Spring before the germination of annual grass weeds listed. Water after application to wash chemical into the turf. SPREADER SETTING: CYCLONE 3.

PRECAUTION: Do not apply in the Spring to completely new lawns started the previous Fall. Do not apply to bentgrass putting greens or dichondra turf. Do not apply to newly sprigged areas of Bermudagrass, St. Augustinegrass or Centipedegrass until these turfs are well established since root inhibition may result. Do not reseed within 42 days after treatment. ACTIVE INGREDIENT:

> N-butyl N-ethyl-a,a,a-trifluoro-2,6-dinitr-p-toluidine INERT INGREDIENTS:

2.5 97.5 *BALAN the registered trademark for benefin, Elanco Products Co.

CAUTION: KEEP OUT OF REACH OF CHILDREN

Harmful if swallowed or absorbed through the skin. Causes eye irritation. Avoid contact with skin, eyes or clothing. In case of contact flush with water. Wash thoroughly with soap and water after use. Do not contaminate foodstuffs or feed.

Do not feed forage from treated areas to livestock. Destroy bag when empty. Do not reuse. This product is toxic to fish. Keep out of lakes, streams or ponds. Do not apply where runoff is likely to occur. Do not contaminate any body of water by direct application, cleaning of equipment or disposal of wastes.

Net Weight 25 lbs.

Manufactured for

Lawn * Doctor... Wickatunk, New Jersey

ROCKLAND (Management -LU., IN**C**,

P 6. 5. X 204

CALDWELL, N. J. 07006