

[572-279]

Net Contents:

ACCEPTED
JUL 3 1974
UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND ROENTGENIC ACT
FOR ECONOMIC POISON REGISTERED
EPA UNDER NO. 572-278



SUPER "KILL" INSECTICIDE SPRAY

FOR MILK ROOMS, DAIRY BARN, CATTLE, CALVES, HOGS AND PIGS AND FEED AND GRAIN STORAGE AREAS.

IBA SUPER "KILL" INSECTICIDE SPRAY contains a high percentage of Pyrethrin and Piperonyl Butoxide in a light mineral oil base and is formulated for use in most any type fogging machine that will produce a very fine mist spray for thermal fog.

DIRECTIONS FOR USE

IBA SUPER "KILL" INSECTICIDE SPRAY is ready to use. Use as directed to kill stable flies, horn flies, house flies, mosquitoes, gnats and exposed stages of cereal insects.

BARN SPRAY: Use in barns, milk houses, out-buildings, pens for effective control of all flies. Spray on walls, ceilings, pens, posts, stalls, windows, doors, where flies congregate. Close doors and windows for 10 minutes when spraying to avoid draft, waste and obtain better results.

Use fogger, power or compressed air sprayer, if preferred, that throws a good fine even spray. Direct the mist toward the upper portion of the enclosure filling the air with fog or fine mist, using 2 ounces of spray per 1000 cubic feet of air space.

ANIMAL SPRAY — CATTLE, CALVES, HOGS: Spray animals with a fine mist at least once a day to kill horn flies, houseflies, mosquitoes.
DAIRY CATTLE — for killing these flies, spray at milking time and only when necessary. No more than 2 ounces of spray should be applied

per animal per application. Tabanids to be killed must be hit directly by spray.

TO KILL EXPOSED STAGES OF CEREAL INSECTS, (granary weevil, rice weevil, saw toothed grain beetle, confused flour beetle, rusty grain beetle, red flour beetle, cadelles, meal worms, grain mites, Angoumois grain moth, Indian meal moth and Mediterranean flour moth). Clean and sweep flour and grain bins and chutes. Fog SUPER "KILL" thoroughly with special emphasis to forcing spray into cracks and checks of ceiling, walls and floor. To be effective fog must penetrate hiding places of insects. Infested stored food products should be fumigated or treated by other effective methods using an approved product intended for this purpose.

As a preventative control, fog once a week during warm months, even though insects are not visible.

When insects are visible, repeat fogging as often as necessary.

E.P.A. Est. 572-NJ-1 EPA Reg. No. 572-278A-11403

100% ACTIVE INGREDIENTS

Pyrethrins	0.12%
*Piperonyl Butoxide, Technical	0.48%
Mineral Seal Oil	49.70%
Petroleum Distillate	49.70%
*Equivalent to 0.38% (butylcarbityl) (4-propylpiperonyl) ether and 0.096% related compounds.	

CAUTION: KEEP OUT OF REACH OF CHILDREN

Harmful if swallowed. DO NOT USE, POUR, SPILL, OR STORE NEAR HEAT OR OPEN FLAME. Do not reuse empty containers. Destroy by perforating or crushing. Bury or discard in a safe place away from water supplies. If container is a drum, return to drum reconditioner is an alternative.

Manufactured for

ROCKLAND CHEMICAL CO., INC.
Passaic Ave., N. Caldwell, N.J.