

Page 572 30

P11-25

1/29/71

131

**PRECAUTIONARY STATEMENTS**

**HAZARDS TO HUMANS**

**CAUTION:** Harmful if swallowed. Avoid contact with eyes and skin. Avoid inhalation of spray mist. Avoid contamination of food and feed.

**IF SWALLOWED:** Do not induce vomiting. Call a physician immediately.

**IF ON SKIN:** Wash with soap and water.

**ENVIRONMENTAL HAZARDS**

Do not contaminate water used for irrigation or domestic purposes. Do not apply directly to any body of water. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

**PHYSICAL HAZARDS**

Do not use, pour, spill or store near heat or open flame.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Read entire Directions for Use and warranty before using this product.

Spray weeds and grass in drives, in cracks of walks and patios, around buildings, under fences and other places where weeds and grass are not wanted.

Care must be taken to keep this herbicide away from desirable flower beds, lawns and shrubs. Do not apply over roots of valuable trees or woody ornamental plants. Only apply herbicide outside the drip line of trees. Avoid spray drift to desirable plants.

Thoroughly wet all foliage to insure contact action. Stand as close to weeds and grass as possible. A jet stream spray is less likely to drift. Best results are obtained when applications are made during or after weed emergence while growth is succulent. One quart will cover 80 square feet or where weeds and grass are tall and dense, 60 square feet. For control of hard to kill perennial weeds and grasses, such as Johnsongrass, Bindweed and Wild Carrot, use one quart to treat 20 square feet.

Although ROCKLAND ready to use WEED & GRASS AWAY has considerable activity through foliar contact, much of its activity is through the roots and, therefore, adequate rainfall to move the chemical into the root zone is necessary.

Under conditions where the rainfall is high and the growing season long, a second application may be necessary to maintain a weed free soil surface.

**STORAGE AND DISPOSAL**

**STORAGE:** Store unused material in original container only in a locked storage area out of reach of children and animals. Keep container tightly closed. Store at temperature above 32°F.

**DISPOSAL:** Do not use empty container. Securely wrap in several layers of newspaper and discard in trash. Rinse thoroughly before discarding in trash.

**WARRANTY:** Seller makes no warranty, expressed or implied, concerning the use of this product other than that specified on the label. Buyer assumes all risk of use and/or handling of this material when he uses and/or handles in conformity to label instructions.

Manufactured by

 **Rockland Chemical Co., Inc.**  
West Caldwell, N.J. 07068 U.S.A.

Product # 308125

EPA Reg. No. 572-308

EPA Est. 572-NJ-2

© Rockland Chemical Co., Inc. 1963

# Rockland<sup>®</sup> Weed and Grass Away



- Ready-to-use
- Kills weeds and grass in walks, drives, patios

**ACTIVE INGREDIENTS:**

Prometon; 2,4-bis (isopropylamino)-6-methoxy-  
s-triazine

1.5%

**INERT INGREDIENTS:**

98.5%

**CAUTION: KEEP OUT OF REACH OF CHILDREN**  
See back panel for additional cautions.  
Net Contents One Quart (32 Fl. Oz.)

Not Reviewed. Registrant claims to be in accordance with Reregistration of Pesticide Product Guidance for Phase 2 Response, Page 2.12, Active to Inert Change in Status.

**BEST AVAILABLE COPY**

00 20 07

W 0 1 3

hogs and for fly control around stables. Also for use in milk rooms, dairy barns, feed and grain storage bins and fly control on dairy and beef cattle and hogs.

**ACTIVE INGREDIENTS:**

Pyrethrins.....0.12%  
\*Piperonyl Butoxide, Tech.....0.48%

\*INERT INGREDIENTS.....99.40%  
100.00%

\*Equivalent to 0.36% (butylcarbityl, 6-propylpiperonyl) ether and 0.096% related compounds.

\* \* CONTAINS AROMATIC PETROLEUM SOLVENT  
**KEEP OUT OF REACH OF CHILDREN**

**CAUTION**

See side panel for additional precautionary statements.

**NET CONTENTS 1 GALLON**

**BEST AVAILABLE COPY**

**Not Reviewed. Registrant claims to be in accordance with Reregistration of Pesticides Product Guidance for Phase 2 Response, Page 2.12, Active to Inert Change in Status.**

**PRECAUTIONARY STATEMENTS**

**HAZARDS TO HUMANS AND DOMESTIC ANIMALS: CAUTION -** Harmful if swallowed. Avoid contact with skin, eyes or clothing. Avoid breathing spray mist. Avoid contamination of food, feed and watering troughs. Do not fog very young animals. Wash after handling.

**ENVIRONMENTAL HAZARDS:** This product is toxic to fish. Do not apply directly to water or wetlands. Do not contaminate water by cleaning of equipment or disposal of waste.

**PHYSICAL OR CHEMICAL HAZARD:** Do not use or store near heat or open flame.

**STATEMENT OF PRACTICAL TREATMENT**

**IF SWALLOWED:** Do not induce vomiting. Petroleum distillate may cause aspiration pneumonia. Call physician or Poison Control Center immediately.

**IF IN EYES:** Flush with plenty of water. Get medical attention if irritation persists.

**IF INHALED:** Remove to uncontaminated atmosphere.

**IF ON SKIN:** Remove contaminated clothing and wash affected areas with soap and water. Launder clothing before reuse.

**STORAGE AND DISPOSAL**

Do not contaminate water, food or feed by storage or disposal.

**STORAGE:** Store unused material in original container only in a locked, dry storage area out of reach of children and animals. Keep container tightly closed.

**PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility in accordance with state and local procedures.

**CONTAINER DISPOSAL:** One Gallon. Do not reuse empty container. Securely wrap in several layers of newspaper and discard in trash. Rinse thoroughly before discarding in trash. Five Gallons and larger - Three rinse for equivalent. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill or by other procedures approved by state and local authorities.

**WARRANTY:** Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

EPA Reg. No. 572-278

EPA Est. No. 572-101

Manufactured by

**Rockland Chemical Co., Inc.**

West Caldwell, NJ 07006 USA

02789

982

**DIRECTIONS FOR USE**

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. Livestock fogging spray is ready to use. Use as directed to kill stable flies, horn flies, house flies, mosquitoes, gnats and exposed stages of cereal insects.

**HORSES AND PONIES:** Spray insect spray from the front of the animal with the direction of the hair, making certain spray nozzle is at least 5 feet from animal. Use not more than an ounce per day. Do not apply to horses intended for human consumption.

**BARN & STABLE SPRAY:** Use in barns, milk houses, outbuildings and pens for effective control of all flies. Spray on walls, ceilings, pens, posts, stalls, windows and doors, where flies congregate. Close doors and windows for 10 minutes when spraying to avoid draft, waste and obtain better results. Use fogger, power or compressed air sprayer if preferred, that throws a good, fine even spray. Direct the mist toward the upper portion of the enclosure filling the air with fog or fine mist, using 2 ounces of spray per 1000 cubic feet of air space.

**CATTLE, CALVES, HOGS:** Spray animals with a fine mist at least once a day to kill horn flies, houseflies, mosquitoes.

**DAIRY CATTLE:** For killing these flies, spray at milking time and only when necessary. No more than 2 ounces of spray should be applied per animal per application. Animals to be killed must be hit directly by spray.

**TO KILL EXPOSED STAGES OF CEREAL INSECTS:** Granary weevil, rice weevil, saw toothed grain beetle, confused flour beetle, rusty grain beetle, red flour beetle, larvae, meal worms, grain mites, Angoumois grain moth, Indian meal moth and Mediterranean flour moth. Clean and sweep flour and grain bins and chutes. Fog Horse, Dairy and Livestock Insect Spray thoroughly with special emphasis to forend spray into cracks and checks of ceiling, walls and floor. To be effective, fog must penetrate hiding places of insects. Infested stored food products should be fumigated or treated by other effective methods using an approved product intended for this purpose.

As a preventative control, fog once a week during warm months, even though insects are not visible.

When insects are visible, repeat fogging as often as necessary.

