

excessive runoff. Treat when pests appear and repeat at 7 to 10 day intervals, if needed. Note: Do NOT use on azaleas, camelias, poinsettias, rose bushes or variegated ivy because of possible injury to these plants.

Dosage and Dilutions For Use On Ornamental Plants

Pests Controlled	Amount of DURSBAN 2-E in Water to Make		
	3 gallons	10 gallons	100 gallons
Bagworms			
Grasshoppers			
Mites	0.5 fl. oz.	1.6 fl. oz.	1 pint*
Spittlebugs			
Whiteflies			
Cutworms			
Mealybugs	1 fl. oz.	3.2 fl. oz.	1 quart**
Leafhoppers			
Thrips (exposed)			

*Equivalent to 4 ounces of chlorpyrifos per 100 gal.

**Equivalent to 8 ounces of chlorpyrifos per 100 gal.

Note: Mosquitoes coming to rest on areas treated for control of turf and ornamental plant pests will be killed for varying periods of time after treatment, depending on exposure of treated areas to weathering conditions.

WARNING

KEEP OUT OF REACH OF CHILDREN

May be fatal if swallowed. May be absorbed through skin. May be injurious to eyes and skin. Combustible

Do not take internally. Do not get in eyes or on skin.

Wash thoroughly after handling. Wash contaminated clothing before re-use. Flush contaminated eyes with plenty of water and get prompt medical attention. Avoid breathing vapors and spray mist. Avoid spraying food crops. Keep away from food, feedstuffs and domestic water supplies. Keep container closed. Do not use, pour, spill or store near heat or open flame. Handle concentrate in a ventilated area.

Note to Physician: Chlorpyrifos is a cholinesterase inhibitor. Treat symptomatically. Atropine is an antidote. Keep children and pets off treated areas until spray has dried. Keep out of lawn and garden fishpools and other bodies of water. Fish may be killed if their waters are contaminated with this product. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated.

This product is toxic to fish, birds and other wildlife. Keep out of lakes, streams or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

Notice: Seller warrants that the product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use but neither this warranty nor any other warranty of MERCHANTABILITY or FITNESS FOR A PARTICULAR PURPOSE, express or implied, extends to the use of this product contrary to label instructions or under abnormal conditions, or under conditions not reasonably foreseeable to seller; and buyer assumes the risk of any such use.

© 1984, FMC Technologies, Inc. All rights reserved.

Cont

TURF
1975
 Insecticide
 Ants, Earwigs, Grasshoppers, Sod Webworms
 12-256

PESTS OUT-OF-DOORS: General information: Use DURSBAN 2-E for control of certain pests of lawns and ornamental plants. Consult the following directions for dosage and application details to control the pests listed.

DIRECTIONS FOR USE

Pests of Lawns and other Ornamental and Recreational Turf Grass Areas: Use DURSBAN 2-E insecticide to control Ants, Brown Dog Ticks, Chinch Bugs, Crickets, Earwigs, Grasshoppers and Sod Webworms (lawn moths). Mix DURSBAN 2-E with water according to directions given in the following table and apply as a coarse low pressure spray using suitable application equipment. For best results, the lawn should be moist at time of treatment and care should be taken to obtain complete and uniform coverage of pest infested areas. Spray when pests first appear, repeat when needed.

Dosage and Dilutions For Use On Turf

Areas to be Treated	Amount of DURSBAN 2-E	Dilute with Water to Give
500 sq. ft.	3/4 fl. oz.	15 gallons
1,000 sq. ft.	1 1/2 fl. oz.	30 gallons
5,000 sq. ft.	7 1/2 fl. oz.	150 gallons
10,700 sq. ft.	1 pint	325 gallons
One Acre	2 quarts	1,300 gallons

To control certain white grubs (larvae of European chafer and Japanese beetle) use two to four times the amounts of DURSBAN 2-E shown in the TURF table, i.e., 3 to 6 fl. oz. per 1,000 sq. ft. or 1 to 2 gallons

per acre. Apply in enough water to obtain uniform coverage of the area treated followed immediately by thorough watering to wash the insecticide into the turf. For best results treat when grubs are young and actively feeding near the soil surface, usually during late July and August. Consult your local Agricultural Extension Service specialist for proper time to treat in your area.

To control Turfgrass (Hyperodes) Weevils, use twice the amount of DURSBAN 2-E shown in the above turf table, i.e., 1 gallon diluted in 1,300 gallons of water per acre. Spray suspected problem areas in mid-April and again in mid-May. Consult your local Agricultural Extension Service specialist for proper time to treat in your area.

Note: To help prevent infestation of buildings by Ants, Crickets, Earwigs and Sowbugs (Pillbugs), treat a band of soil 6 to 10 feet wide around and adjacent to the building, also the building foundation to a height of 2 to 3 feet, where pests are active and may find entrance. Use the same dosage and dilution recommended for control of these pests on turf as shown in the above table. Apply as a coarse spray to thoroughly and uniformly wet the band area.

Pests of Ornamentals: Use DURSBAN 2-E to control Bagworms, Cutworms, Grasshoppers, Leafhoppers, Mealybugs, Mites, Spittlebugs, Thrips (exposed) and Whiteflies on flowers, shrubs, vines, shade and flowering trees and evergreens found to be infested with these pests. Dilute DURSBAN 2-E with water according to directions given in the following table and apply using suitable hand or power spray equipment in a manner to provide complete and uniform coverage of infested plants. Avoid over-spraying to the point of