



ROCKLAND CHEMICAL CO., INC.

P. O. BOX 809
WEST CALDWELL, N. J. 07006

739

Rockland (R)

GENERAL PURPOSE AQUEOUS INSECTICIDE

Indoor-outdoor spray
Ready-to-use insecticide
Kills spiders, ants, roaches, fleas
wasps and others listed

JUL 20 1984
572-287

Active Ingredients:

Pyrethrins.....	0.1%
*Technical Piperonyl Butoxide.....	1.0%
Petroleum distillate.....	0.4%
Inert Ingredients:.....	98.5%
	<u>100.0%</u>

*Equivalent to 0.8% (butylcarbityl) (6-propylpiperonyl) ether and to 0.2% related compounds

(15 point type) CAUTION: KEEP OUT OF REACH OF CHILDREN *(12 point type)*
See back panel for additional cautions

Net Contents ONE GALLON

EPA: Front panel caution statements will meet type size requirements according to label size needed for various containers. STORAGE AND DISPOSAL title will meet relative type size requirements.

BEST DOCUMENT AVAILABLE

ROCKLAND CHEMICAL CO., INC.

P. O. BOX 899

WEST CALDWELL, N. J. 07006

PRECAUTIONARY STATEMENTS

CAUTION: Harmful if swallowed. Avoid contamination of feed and food-stuffs. Remove pets, birds and cover fish aquaria before spraying.

ENVIRONMENTAL CAUTIONS

This product is toxic to fish. Do not apply directly to water. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labelling.

Effective, fast acting insecticide for Horses, Dairy Animals, Livestock and for use in Homes, Institutions, Restaurants, Food Processing Plants, Dairies, Livestock and outdoors for Mosquito control. Leaves no oily or objectionable residue.

INDOOR APPLICATION: The use of this product in food processing establishments should be confined to time periods when the plant is not in operation. Food should be removed or covered during treatment. All food processing surfaces should be covered during treatment or thoroughly cleaned before using. Vacate treated area and ventilate before re-occupying. For maximum effectiveness, a combination of spot surface treatment and fogging is recommended.

Apply with conventional mechanical or compressed air equipment.

SURFACE SPRAYING: To control Roaches, Ants, Silverfish, spiders, Crickets, Clover Mites, Cheese Mites, Granary Weevils, Rice Weevils, Confused Flour Beetles, Saw-toothed Grain Beetles, Spider Beetles, Cigarette Beetles, Drugstore Beetles, Meal Worms, Grain Mites, and Cadelles: Adjust sprayer to deliver a coarse wet spray. Direct the spray into hiding places, cracks and crevices, under pallets, around containers of stored foods, around the base of machinery, behind shelves and drawers. For Silverfish, spray bookcases. For Ants, spray trails, nests and points of entry. If surface application only is to be used, spray floors, walls, and other surfaces applying at a rate of 1 gallon to 750 square feet of surface.

To control Carpet Beetles: Spray edges of carpeting and under carpeting and rugs. Make localized application to floor and baseboards. Spray directly into cracks, closets and infested areas of shelving. Repeat treatment as needed.

To control Bedbugs: Spray mattresses lightly, particularly around tufts and seams. Take beds apart and spray in all joints. Treat baseboards, mouldings and floors. Repeat treatment as needed.

To control Fleas and Brown Dog Ticks: Thoroughly spray infested areas, pet beds, resting quarters, nearby cracks and crevices, along and behind baseboards, mouldings, window and door frames, and localized areas of floor and floor covering. Fresh bedding should be placed in animal quarters following treatment. Concurrent treatment of animals with an approved insecticide is recommended. Repeat treatment as needed.

BEST DOCUMENT AVAILABLE

ROCKLAND CHEMICAL CO., INC.

P. O. BOX 719

WEST CALDWELL, N. J. 07006

To control Clothes Moths: Remove any infested articles from storage, brush thoroughly and air for several hours in sunlight, if possible. Apply spray liberally to empty chests, closets, bureaus and other storages, directing the spray into cracks, joints and crevices. After airing, infested articles may be treated lightly keeping nozzle at least three feet from fabric to avoid staining.

ON ANIMALS: To protect cattle and horses from Horn Flies, House Flies, Mosquitoes, and Gnats, apply a light mist sufficient to wet the surface of the hair. To control Stable Flies, Horse Flies and Deer Flies, apply at a rate of 2 ounces per adult cow or horse, sufficient to wet the hair thoroughly. Repeat treatment daily or at intervals necessary to give continued protection.

To control Blood-sucking Lice, apply to the infested areas of the animal, using a stiff brush to get the spray to the base of the hair. Repeat every 2 or 3 weeks if required.

SPACE SPRAYING: Close doors and windows and shut off ventilating systems. Apply with a good machine adjusted to deliver a fine mist or fog. For rapid control of Flies, Fruit Flies, Gnats, Mosquitoes, Skipper Flies, Wasps, Hornets, Angoumois Grain Moths, Mediterranean Flour Moths, Indian Meal Moths, and Tobacco Moths: direct spray at an upward angle, distributing it uniformly throughout the entire area, at a rate of 3-6 ounces per 1000 cubic feet of space.

When animals are present in sprayed buildings, after five (5) minutes, restart ventilating system or remove animals.

OUTDOOR ADULT MOSQUITO CONTROL: Apply as a fine mist to thoroughly treat the entire area. For best results, treat when the wind velocity does not exceed 5 miles per hour. Treat shrubbery or vegetation where mosquitoes may rest. Shrubbery and vegetation around stagnant pools, marshy areas, ponds and shore may be treated. Apply at a rate of 1½ gallons per acre (4½ ounces per 1000 square feet).

APPLICATION OF THIS PRODUCT TO ANY BODY OF WATER IS NOT ACCEPTABLE.

Using truck-mounted conventional misting equipment such as Rotomist® models, 51C, 91 or 100 HT. traveling at 5 m.p.h., to treat a 300 foot swath, calibrate equipment to deliver 4.5 gallons per minute.

STORAGE AND DISPOSAL

STORAGE: Store unused material in original container in a locked storage area out of reach of children and animals. Keep container tightly closed.

DISPOSAL: Do not contaminate water, food or feed by storage or disposal. Triple rinse container (or equivalent). Then dispose of in a sanitary landfill or by other approved State and local procedures. Puncture metal containers. Rinsate that cannot be used should be disposed of in a landfill approved for pesticides or buried in a safe place away from water supplies.

BEST DOCUMENT AVAILABLE

ROCKLAND CHEMICAL CO., INC.

P. O. BOX 809

WEST CALDWELL, N. J. 07006

WARRANTY: Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

Manufactured by

Rockland Chemical Co., Inc.
West Caldwell, NJ U.S.A.

Product

EPA Reg. No. 572-287

EPA Est. 572-NJ-1

BEST DOCUMENT AVAILABLE