


Rockland[®] Horse and Livestock Fly Spray



PM 17 572-287
1-24-96

Not Reviewed. Registrant claims to be in accordance with
 Reregistration of Pesticide Product Guidance for Phase 2
 Response, Page 2.12, Active to Inert Change in Status.

ACTIVE INGREDIENTS:
 Pyrethrins.....0.1%
 Piperonyl Butoxide, Tech.....1.0%
*** INERT INGREDIENTS.....98.9%**

*Equivalent to 0.3% (butylcarbityl)(6-propylpiperonyl) ether and to 0.2% related compounds.
 **Contains Petroleum Distillate.

KEEP OUT OF REACH OF CHILDREN
CAUTION

Rockland Horse & Livestock Fly Spray is an Effective, Fast Acting Insecticide for Horses, Dairy Animals and Livestock.

PRECAUTIONARY STATEMENTS

Hazards to Human and Domestic Animals: CAUTION. Harmful if swallowed. Avoid contamination of feed and foodstuffs. Avoid contact with skin and eyes. Remove pets, birds and cover fish aquariums before spraying. **Environmental Hazards:** This product is toxic to fish. Keep out of lakes, streams or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from area treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

STATEMENT OF PRACTICAL TREATMENT


If Swallowed: Do not induce vomiting. Call physician or Poison Control Center immediately. **If In Eyes:** Flush with plenty of water. Get medical attention. **If On Skin:** Remove contaminated clothing. Wash affected area with soap and water. Launder clothing before reuse. **If Inhaled:** Remove from contaminated atmosphere.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. To protect **Cattle** and **Horses** from Horn Flies, House Flies, Mosquitoes, and Gnats, apply a light mist sufficient to wet the surface of the hair. To control Stable Flies, Horse Flies and Deer Flies, apply at a rate of 2 ounces per adult cow or horse, sufficient to wet the hair thoroughly. Repeat treatment daily or at intervals necessary to give continued protection. To control Blood-Sucking lice, apply to the infested areas of the animal, using a stiff brush to get the spray to the base of the hair. Repeat every 2 or 3 weeks if required.

STORAGE AND DISPOSAL

Storage: Store unused material, in original, closed container only, in a locked storage area out of reach of children and animals. Do not contaminate water, food or feed by storage or disposal. **Disposal:** Do not reuse empty container. Securely wrap in several layers of newspaper and discard in trash.



Rockland Rockland Chemical Co., Inc.
West Caldwell, N.J. 07008 USA