

572-287

6/12/2002

1/4

Rockland® Pyrenone General Aqueous Insecticide

Alt Brand Name: **Horse and Livestock Fly Spray**

- Water Base Product
- Leaves No Oily or Objectionable Residue.
- Effective Fast Acting Insecticide.
- For Use in Homes, Institutions, Restaurants, Feed Processing Plants, Dairies, Livestock and Outdoors for Mosquito Control and for Control of Fleas and Brown Dog Ticks on Premises.
- Kills Crawling & Flying Insects on Contact.
- Contains Natural (Pyrethrin) Insecticide Derived from Flowers.
- House & Garden Insect Killer.
- Plant Spray Bug Killer.

ACTIVE INGREDIENTS:

Pyrethrins.....0.1%
 *Piperonyl Butoxide, technical 1.0%
 INERT INGREDIENTS:.....98.9%


*Equivalent to 0.8% (butylcarbityl) (6-propylpiperonyl) ether and to 0.2% related compounds.

KEEP OUT OF REACH OF CHILDREN

CAUTION

See Back Panel for complete precautionary statements

Net Contents: One Quart, One Gallon



PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS: CAUTION: Harmful if swallowed. Avoid contamination of feed and foodstuffs. Avoid contact with skin and eyes. Remove pets, birds and cover fish aquariums before spraying. **ENVIRONMENTAL HAZARDS:** This product is toxic to fish. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from area treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

FIRST AID	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment	
If in Eyes:	-Hold eye open and rinse slowly and gently with water for 15-20 minutes. -Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. -Call a poison control center or doctor for treatment advice.
If Swallowed:	-Call poison control center or doctor immediately for treatment advice. -Have person sip a glass of water if able to swallow. -Do not induce vomiting unless told to do so by the poison control center or doctor. -Do not give anything by mouth to an unconscious person.
If on Skin or Clothing:	-Take off contaminated clothing. -Rinse skin immediately with plenty of water for 15-20 minutes. -Call a poison control center or doctor for treatment advice.
If inhaled:	-Move person to fresh air. -If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. -Call a poison control center or doctor for further treatment advice.
For information regarding medical emergencies or pesticide incidents, call the International Poison Center at 1-888-740-8712.	

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

FARM DIRECTIONS

Horse & Livestock Fly Spray is an effective, fast acting insecticide for Horses, Dairy Animals and Livestock.

To protect Cattle and Horses from Horn Flies, House Flies, Mosquitoes, and Gnats, apply a light mist sufficient to wet the surface of the hair. To control Stable Flies, Horse Flies and Deer Flies, apply at a rate of 2 ounces per adult cow or horse, sufficient to wet the hair thoroughly. Repeat treatment daily or at intervals necessary to give continued protection.

To Control Blood-Sucking lice, apply to the infested areas of the animal, using a stiff brush to get the spray to the base of the hair. Repeat every 2 to 3 weeks if required.

To Control Stable Flies, Horse Flies and Deer Flies, apply at a rate of 2 ounces per adult animal sufficient to wet the hair thoroughly. Repeat treatment daily or at intervals necessary to give continued protection.

To Control Poultry Lice, spray roosts, walls and nests or cages thoroughly. This should be followed by spraying over the birds with a fine mist.

For Control of Bedbugs and Mites in Poultry Houses: spray crevices of roost poles,, cracks in walls, and cracks in nests where the bedbugs and mites hide.

INDOOR APPLICATION

the use of this product in food processing establishments should be confined to time periods when the plant is not in operation. Food should be removed or covered during treatment. All food processing surfaces should be covered during treatment. All food processing surfaces should be covered during treatment or thoroughly cleaned with potable water before using.

Vacate treated areas and ventilate before reoccupying.

For maximum effectiveness, a combination of spot, surface treatment and fogging is recommended.

SURFACE SPRAYING: To control **Cockroaches, Palmetto Bugs, Ants, Silverfish, Spiders, Crickets, Clovermites, Cheese Mites, Earwigs, Sowbugs, Boxelder Bugs, Centipedes, Granary Weevils, Rice Weevils, Confused Flour Beetles, Rust Red Flour Beetles, Sawtoothed Grain Beetles, Spider Beetles, Cigarette Beetles, Drugstore Beetles, Meal Worms, Grain Mites and Cadelles:** Use a good sprayer adjusted to a coarse wet spray. Direct the spray into hiding places, cracks and crevices, under pallets, around containers of stored foods, around the base of Machinery, behind shelves and drawers, behind and beneath sinks, stoves, refrigerators and cabinets, along baseboards, around door and window frames, into dark corners and in and around garbage compactors, garbage chutes and cans. For **Silverfish**, spray bookcases. For **Ants**, spray trails, nests and points of entry. If surface application only is to be used, spray floors and other surfaces applying at a rate of 1 gallon to 750 square feet of surface.

In case of infested foods in cabinets, dispose of the infested material. Remove all contents and clean thoroughly. Spray interior surfaces of empty cabinets thoroughly. After spray has dried, cover shelves with paper or otherwise protect foods from direct contact with sprayed surfaces.

To Control Carpet Beetles: Spray edges of carpeting and under carpeting and rugs. Make localized application to floor and baseboards. Spray directly into cracks, crevices and infested areas of shelving. Repeat treatment as needed.

To Control Bedbugs: Spray mattresses lightly, particularly around tufts and seems. Take beds apart and spray in all joints., Treat baseboards, moldings and floors. Repeat treatment as needed.

Pet Areas: To control Fleas and Brown Dog Ticks on premises: Thoroughly spray infested areas, pet beds, resting quarters, nearby cracks and crevices, along and behind baseboards, moldings, window and door frames, and entire areas of floor and floor covering. Fresh bedding should be placed in animal quarters following treatment. Concurrent treatment of animals is recommended. Repeat treatment as needed. Do not apply directly to household pets.

To Control Clothes Moths: Remove any infested articles from storage, brush thoroughly and air for several hours in sunlight, if possible. Apply spray liberally to empty chests, closets, bureaus and other storages directing the spray into cracks, joints and crevices. After airing, infested articles may be treated lightly keeping nozzle at least three feet from fabric to avoid staining.

Space Spraying: Close doors, windows and transoms and shut off ventilating systems. Apply with a good machine adjusted to deliver a fine mist or fog. For rapid control of **House Flies, Fruit Flies, Gnats, Mosquitoes, Skipper Flies, Wasps, Hornets, Angoumois Grain Moths, Mediterranean Flour Moths, Indianmeal Moths and Tobacco Moths**, direct spray at an upward angle distributing it uniformly throughout the entire area at a rate of 3-6 ounces per 1000 cubic feet of space. Do not remain in treated areas. If insects have not dropped to floor in 3 minutes, repeat spraying as quantity sprayed was insufficient. Ventilate the area after treatment is completed. To aid in the prevention of breeding, spray on standing water and in garbage pails.

Fogging: To kill exposed or accessible stages of the above named crawling insects, use a rate of 1 gallon to 20,000 to 40,000 cubic feet of space. Contact insects directly whenever possible.

4/4₄

Adult Mosquito Control: Apply as a fine mist to thoroughly treat the entire area. For best results, treat when the wind velocity does not exceed 5 miles per hour. Treat shrubbery or vegetation where Mosquitoes may rest (low, marshy places, stagnant pools or ponds and shore lines). Apply at a rate of 2.5 pints per acre (0.1 ounce per 100 sq. ft.). Using truck mounted conventional misting equipment such as Rotomist® models 51C, 91 or 100HT traveling at 5 m.p.h. to treat 300 ft. swath, calibrate equipment to deliver 0.9 gallon per minute.

**(ALTERNATE) DIRECTIONS FOR USE (CONSUMER LABELING)
SHAKE WELL BEFORE USE.**

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

TO KILL---

--Crawling Insects: Cockroaches, Silverfish, Sowbugs, Ants, Crickets, Earwigs, Grain Mites, Clovermites, Spiders, Boxelder Bugs, Centipedes. Turn sprayer nozzle to stream position and spray directly into cracks, crevices and other hiding places along baseboards, hitting as many bugs as possible. Repeat as necessary. Keep about three feet away from walls, fabrics and furniture.
Flying Insects: Flies, Houseflies, Mosquitoes, Gnats, Mediterranean Flour Moths, Small Flying Moths. Turn sprayer nozzle to spray position. Keep about three feet away from walls, fabrics and furniture.

--Wasps & Hornets: Turn sprayer nozzle to stream position. Spray directly on insects whenever possible. Apply a wet spray after dark when all insects have returned to their nests.

-- Garden Insects on House Plants and Ornamental Plants Outdoors: Turn spray nozzle to spray position. Direct spray so that upper and lower leaf surfaces are contacted. To control **Aphids, Bagworms, Thrips, Flea Beetles, Japanese Beetles, Lace Bugs, Leaf Hoppers, Leaf tiers, Mimosa Webworm Larvae, Plant Bugs and Whiteflies** use fine mist spray about 18 to 24 inches from infested plants. Use caution on new growth and tender foliage. Repeat treatment as necessary. Avoid spraying plants when temperature exceeds 85° F. Spray when air is still. Avoid heavy wetting.

--Pet Areas: To control **Fleas and Brown Dog Ticks** on premises: Turn sprayer nozzle to spray position. Thoroughly spray infested areas, pet beds, resting quarters, nearby cracks and crevices, along and behind baseboards, moldings, window and door frames, and entire areas of floor and floor covering. Fresh bedding should be placed in animal quarters following treatment. Concurrent treatment of animals is recommended. **DO NOT USE** this product on Household Pets.

STORAGE AND DISPOSAL

STORAGE: Store unused material, in original, closed container only, in a locked storage area out of reach of children and animals. Do not contaminate water, food or feed by storage or disposal.

DISPOSAL: If Empty: Do not reuse this container. Place in trash or offer for recycling if available.

If Partly Filled: Call your local solid waste agency or 1-800-CLEANUP for disposal instructions.

Never place unused product down any indoor or outdoor drain.

WARRANTY: Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

EPA Est. No. 572-NJ-1
EPA Reg. No. 572-287
Product C28732,01
0402

UPC

Manufactured by
Rockland Corporation
West Caldwell, NJ 07006