

ROCKLAND

AC

APR 09 1975

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 6-12-222

TREATS 12,500 SQ. FT.

PROFESSIONAL 5% DIAZINON GRANULAR LAWN INSECTICIDE

For Professional Use Only DIRECTIONS FOR USE

OVERALL LAWN APPLICATION

To control the following lawn insects on both grass and dichondra lawns: Chinch Bugs, Earwigs, Ticks, Ants, Lawn Moths (Sod Webworms), Bermuda Mites, Crickets, Armyworms, Brown Dog Ticks, Chiggers, Clover Mites, Cutworms, Fleas, Leafhoppers, Sowbugs, Millipedes, Springtails (Collembola).

Apply to your lawn with a fertilizer spreader at the rate shown in the table below. Application should be made when grass is dry. Water lawn lightly after application. Repeat as necessary. Keep children and pets off treated areas until this material is washed into the soil and grass is dry.

For application with other types of equipment, or by hand, apply at the rate of 2 lbs. per 1000 sq. ft. (an area 20' by 50') or 3.2 oz. per 100 sq. ft. (an area 10' by 10'). For heavy lawns, application should be increased to 3 lbs. per 1000 sq. ft. (or 4.8 oz. per 100 sq. ft.).

Billbugs and White Grubs of Japanese Beetle, European Chafer, Southern Chafer. Apply 2.5 lbs. per 1000 sq. ft. of lawn. Water grass thoroughly after application. For grub control make application any time between late July and early October. For Billbug control, apply when activity is first observed or when chewed or brown grass indicates damage from these insects. Repeat as necessary.

Hyperodes Weevil (New York only). Apply 2 lbs. per 1000 sq. ft. of grass. Water grass thoroughly after application. Treat problem areas in mid-April and again in mid-May.

SPREADER SETTING NO.

(per 1000 sq. ft.)

Insects listed below

Spreader Make and Model	Regular or Dichondra Lawn — 2 lbs.	St. Augustine Lawn — 3 lbs.	White Grubs, Billbugs and Hyperodes Weevil — 2 1/2 lbs.
Scott	5-1 8	6-1 8	5-5 8
Jackson 18-26 (18")	5	5-3 4	5-1 2
Salem Micro Set (22")	9-1 4	8-5 8	8-7 8
Cyclone Models 10, 15, 20	4-3 4	5-3 4	5-1 4
Model B-1	4	5	5-1 2
Lawn Beauty	6	6-7 8	6-1 2
Greenfield Model 103	4	5	5-1 2
Model 105	7	7-1 2	8

Chinch Bugs, Earwigs, Ants, Bermuda Mites, Lawn Moths (Sod Webworms), Crickets, Armyworms, Brown Dog Ticks, Chiggers, Clover Mites, Cutworms, Fleas, Leafhoppers, Sowbugs, Millipedes, Springtails (Collembola).

These suggestions are for your guidance in spreading this material.

SPOT TREATMENT

ANTS: Apply 1-1 2 teaspoons over and around each individual ant hill.

DIGGER WASPS: Apply 1 1/2 teaspoons in and around each individual ground nest opening. Make the application about dusk when the wasps are not active.

BAND TREATMENT AROUND FOUNDATION

The following insects are commonly found around or near the foundations of houses as well as in the lawn. Certain of these insects may even enter the house.

Ants, Brown Dog Ticks, Clover Mites, Crickets, Earwigs, Fleas, Millipedes, Sowbugs, Springtails (Collembola).

For control of the above insects, apply a five-foot band next to the foundation of the house. Distribute the granules uniformly using a rate of 3.2 oz. for each 5' by 20' area to be treated.

CAUTION: Keep out of reach of children. Harmful if swallowed. Avoid contamination of food and feed. Do not store near food or feed products. Avoid inhalation of dust. Avoid contact with skin, eyes and clothing. Wash thoroughly after handling.

LAWN INSECTICIDE

For Professional Use Only

DIRECTIONS FOR USE

OVERALL LAWN APPLICATION

To control the following lawn insects on both grass and dichondra lawns: Chinch Bugs, Earwigs, Ticks, Ants, Lawn Moths (Sod Webworms), Bermuda Mites, Crickets, Armyworms, Brown Dog Ticks, Chiggers, Clover Mites, Cutworms, Fleas, Leathoppers, Sowbugs, Millipedes, Springtails (Collembola).

Apply to your lawn with a fertilizer spreader at the rate shown in the table below. Application should be made when grass is dry. Water lawn lightly after application. Repeat as necessary. Keep children and pets off treated areas until this material is washed into the soil and grass is dry.

For application with other types of equipment, or by hand, apply at the rate of 2 lbs. per 1000 sq. ft. (an area 20' by 50') or 3.2 oz. per 100 sq. ft. (an area 10' by 10'). For heavy lawns, application should be increased to 3 lbs. per 1000 sq. ft. (or 4.8 oz. per 100 sq. ft.).

Billbugs and White Grubs of Japanese Beetle, European Chafer, Southern Chafer. Apply 2.5 lbs. per 1000 sq. ft. of lawn. Water grass thoroughly after application. For grub control make application any time between late July and early October. For Billbug control, apply when activity is first observed or when chewed or brown grass indicates damage from these insects. Repeat as necessary.

Hyperodes Weevil (New York only). Apply 2 lbs. per 1000 sq. ft. of grass. Water grass thoroughly after application. Treat problem areas in mid-April and again in mid-May.

SPREADER SETTING NO.

(per 1000 sq. ft.)

Insects listed below

Spreader Make and Model	Regular or Dichondra Lawn — 2 lbs.	St. Augustine Lawn — 3 lbs.	White Grubs, Billbugs and Hyperodes Weevil — 2 lbs.
Scott	5-1/8	6-1/8	5-5/8
Jackson 18-26 (18")	5	5-3/4	5-1/2
Salem Micro Set (22")	9-1/4	8-5/8	8-7/8
Cyclone Models 10, 15, 20	4-3/4	5-3/4	5-1/4
Model B-1	4	5	5-1/2
Lawn Beauty	6	6-7/8	6-1/2
Greenfield Model 103	4	5	5-1/2
Model 105	7	7-1/2	8

Chinch Bugs, Earwigs, Ants, Bermuda Mites, Lawn Moths (Sod Webworms), Crickets, Armyworms, Brown Dog Ticks, Chiggers, Clover Mites, Cutworms, Fleas, Leathoppers, Sowbugs, Millipedes, Springtails (Collembola).

These suggestions are for your guidance in spreading this material.

SPOT TREATMENT

ANTS: Apply 1-2 teaspoons over and around each individual ant hill.

DIGGER WASPS: Apply 1-2 teaspoons in and around each individual ground nest opening. Make the application about dusk when the wasps are not active.

BAND TREATMENT AROUND FOUNDATION

The following insects are commonly found around or near the foundations of houses as well as in the lawn. Certain of these insects may even enter the house.

Ants, Brown Dog Ticks, Clover Mites, Crickets, Earwigs, Fleas, Millipedes, Sowbugs, Springtails (Collembola).

For control of the above insects, apply a five-foot band next to the foundation of the house. Distribute the granules uniformly using a rate of 3.2 oz. for each 5' by 20' area to be treated.

CAUTION: Keep out of reach of children. Harmful if swallowed. Avoid contamination of food and feed. Do not store near food or feed products. Avoid inhalation of dust. Avoid contact with skin, eyes and clothing. Wash thoroughly after handling. May be absorbed through skin.

FISH AND WILDLIFE CAUTION: This product is toxic to fish, birds and other wildlife. Keep out of lakes, streams or ponds. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

Do not reuse container. Destroy it by burying or burning. Avoid smoke or fumes.

CAUTION: KEEP OUT OF REACH OF CHILDREN

ACTIVE INGREDIENT

0.0 diethyl 0.2

NET INGREDIENT

MAZINON

Net Weight 25 lbs.

Manufactured by

ROCKLAND CHEMICAL CO.

PASSAIC AVENUE

WEST CALDWELL, NEW JERSEY