



ROCKLAND RABON[®] LIVESTOCK DUST

READY TO USE INSECTICIDE



A Rabon[®] Insecticide Dust for Direct Application to
DAIRY AND BEEF CATTLE

Controls Lice and Horn Flies and aids in the control of Face Flies. Controls Lice on Swine.

CAUTION: KEEP OUT OF REACH OF CHILDREN.

SEE BACK PANEL FOR ADDITIONAL CAUTIONS.

ACTIVE INGREDIENTS:	BY WEIGHT:
2-chloro-1-(2,4,5-trichlorophenyl) vinyl dimethyl phosphate	3.00%
INERT INGREDIENTS:	97.00%
TOTAL	100.00%
*Rabon [®] Insecticide, Shell Chemical Co.	

MANUFACTURED BY

Rockland Chemical Co., Inc.
Passaic Avenue, West Caldwell, N. J.

NET WT. ONE POUND

DIRECTIONS FOR USE

RABON[®] LIVESTOCK DUST is a ready-to-use insecticide dust which can be applied by hand or through the use of self-treating dust bags. There is no withholding period from last application to slaughter.

Animal	Insect	REMARKS
Beef Cattle	Horn Flies	HAND DUSTING—Apply approximately 2 oz. of dust by shaker can, rotary duster or by spoon to the upper portions of the back, neck and poll, and to the face as an aid in the control of face flies. Rub in lightly to carry the dust beneath the hair. Repeat as necessary.
Dairy Cattle	Face Flies	
	Lice	

SELF-TREATING DUST BAG.

FORCED USE — Put RABON[®] LIVESTOCK DUST in cotton cloth or double burlap bags or use pre-packed weather-proof cattle dust bags and hang in barn door exits or alleyways leading from animal buildings, salt or mineral blocks, or watering holes. Protect cloth or burlap bags from weather.

FREE CHOICE USE — Use the same dust bags as above but place in loafing sheds, holding pens, feedlots, near watering holes or other areas where cattle gather. The free choice use aids in the control of lice.

Animal	Insect	REMARKS
Swine	Lice	HAND DUSTING — Apply 3 - 4 oz. of dust by conventional hand or power duster to each animal with special attention given to the neck and around the ears. Repeat as necessary but not more often than once every 14 days. In severe infestations, both individual animals and bedding may be treated. One lb. of 3% dust should be applied per 150 sq. ft. of bedding.

HAND DUSTING — Apply 3 - 4 oz. of dust by conventional hand or power duster to each animal with special attention given to the neck and around the ears. Repeat as necessary but not more often than once every 14 days. In severe infestations, both individual animals and bedding may be treated. One lb. of 3% dust should be applied per 150 sq. ft. of bedding.

CAUTION: Harmful if swallowed. Avoid breathing dust. Do not get in eyes, on skin, or on clothing. If the material gets into the eyes, wash with plenty of water for 15 minutes; if irritation persists, see a physician. Wash thoroughly with soap and water after handling and before eating or smoking. Avoid contamination of feed and foodstuffs. Keep away from heat and open flame. Do not apply in dwellings.

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

Do not reuse empty container. Destroy it by perforating or crushing. Bury or discard in a safe place.

WARRANTY: Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

EPA Reg. No. 577-295

FPA Est. 572-111

OCT. 6 1975

Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, the pesticide is registered under EPA Reg. No. 577-295.

ACCEPTED