

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY

OCT 30 1989

200 25404
17

...
...
...
...

Your attention is directed to the following information:

The label was reviewed to allow, submitted in connection with the registration under the Federal Insecticide, Fungicide, and Rodenticide Act, 1974, for the pesticide, Tetrachlorvinphos, 100% concentrate, to be changed to the label listed below before you release the product for shipment to retail outlets for labeling:

1. Both ACTIVE INGREDIENTS and INERT INGREDIENTS must be aligned to the same margin.
2. Remove outer brackets in active ingredient chemical name and place a colon after Tetrachlorvinphos.
3. Under Environmental Hazards, change "This product is toxic . . ." to "This pesticide is toxic"

55397:1:Hebert:LLR-8:KENCO:10/26/89:12/06/89:DD:EK:aw

CONCURRENCES

SYMBOL							
SURNAME							
DATE							

Submit five (5) copies of your final printed labeling before you release the product for shipment.

A stamped copy of the labeling is enclosed for your records.

Sincerely yours,

George T. LaRocca
Product Manager (15)
Insecticide-Rodenticide Branch
Registration Division (H7505C)

enclosure

Reg no: 572-295

PM-15

3 3 5

ROCKLAND CHEMICAL CO., INC.

P. O. BOX 809

WEST CALDWELL, N. J. 07007-0809

ROCKLAND

RABON^R LIVESTOCK DUST

Ready to Use Insecticide

A Rabon^R Insecticide Dust for Direct Application to DAIRY AND BEEF CATTLE

Controls Lice and Horn Flies and aids in the control of Face Flies. Controls Lice on Swine.

KEEP OUT OF REACH OF CHILDREN

CAUTION

See Back Panel For Additional Precautionary Statements

ACTIVE INGREDIENT

Tetrachlorvinphos [2-chloro-1-(2,4,5-trichlorophenyl) vinyl dimethyl phosphate].....3.0%

INERT INGREDIENTS..... 97.0%
100.0%

EPA Reg. No. 572-295

EPA Est. No. 572-NJ-1


Net Weight one lb

ACCEPTED
with COMMENTS
in EPA Letter Dated:

30 1971

Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No.

572-295



PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS: CAUTION: Harmful if swallowed. Avoid contact with skin and eyes. Avoid breathing dust. Wash thoroughly with soap and water after handling and before eating or smoking. In in eyes, wash with plenty of water. If irritation persists, get medical attention. Wear long-sleeved shirt and pants; chemical resistant gloves; shoes and socks for protection when handling.

NOTE TO PHYSICIAN/VETERINARIAN: This product is a cholinesterase inhibitor. If symptoms of cholinesterase inhibition are present, atropine sulfate, by injection, is antidotal. 2-PAM is also antidotal and may be administered, but only in conjunction with atropine.

ENVIRONMENTAL HAZARDS: This product is toxic to fish. Do not contaminate water when disposing of equipment washwaters.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

ROCKLAND RABON LIVESTOCK DUST is a ready-to-use insecticide dust which can be applied by shaker can or through the use of self-treating dust bags. THERE IS NO WITHHOLDING PERIOD FROM LAST APPLICATION TO SLAUGHTER.

ANIMAL	INSECT	REMARKS
Beef Cattle Dairy Cattle	Horn Flies Face Flies Lice	HAND DUSTING: Apply approximately 2 oz. of dust by shaker can, rotary duster or by spoon to the upper portions of the back, neck and poll, and to the face as an aid in the control of face flies. Rub in lightly to carry the dust beneath the hair. Repeat as necessary.

SELF-TREATING DUST BAG.

FORCED USE — Put ROCKLAND RABON LIVESTOCK DUST in cotton cloth or double burlap bags or use pre-packed weather-proof cattle dust bags and hang in barn door exits or alleyways leading from animal buildings, salt or mineral blocks, or watering holes. Protect cloth or burlap bags from weather.

FREE CHOICE USE — Use the same dust bags as above but place in loafing sheds, holding pens, feedlots, near water holes or other areas where cattle gather. The free choice use aids in the control of lice.

ANIMAL	INSECT	REMARKS
Swine	Lice	HAND DUSTING: Apply 3-4 oz. of dust by conventional hand or power duster to each animal with special attention given to the neck and around the ears. Repeat as necessary but not more than once every 14 days. In severe infestations, both individual animals and bedding may be treated. One lb. of 3% dust should be applied per 150 sq. ft. of bedding.

ROCKLAND RABON LIVESTOCK DUST
P. O. BOX 809
WEST CALDWELL, N. J. 07007-0809

525
Rockland Rabon Livestock Dust

STORAGE: Store unused material, in original container only, in a locked storage area out of reach of children and animals. Keep container tightly closed.

DISPOSAL: Do not reuse empty container. Wrap in several layers of newspaper and discard in trash.

WARRANTY: Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

EPA Reg. No. 572-295
#C29501

EPA Reg. No. 572-N-1-1
1989

NET WEIGHT 1 lb.

