


CRC
ROCKLAND



5/2-35

WEED
&
FEEED "B"

18-6-6

FEEED & "B"

18-6-6

TREATS

5000 SQUARE FEET

- ① Weeds with 2,4-D and Silvex to control most broadleaf lawn weeds.
Feeds with 18-6-6 Fertilizer.**
- ②**
- ③ For Best results mow lawn before applying.**

CAUTION

- ① Weeds with 2,4-D and Silvex to control most broadleaf lawn weeds.
Feeds with 18-6-6 Fertilizer.
- ②
- ③ For Best results mow lawn before applying.

CAUTION

KEEP OUT OF REACH OF CHILDREN

SEE BACK PANEL FOR COMPLETE CAUTION

READ ENTIRE LABEL BEFORE USING THIS PRODUCT

ACTIVE INGREDIENTS:

Dimethylamine Salt of 2,4-Dichlorophenoxyacetic Acid* ...	1.40%
Butoxy propyl ester of Silvex [2-(2,4,5-trichloro- phenoxy) propionic acid]**79%
INERT INGREDIENTS:	97.81%
	100.00%

GUARANTEED ANALYSIS

Total Nitrogen (N)	18%
Available Phosphoric Acid (P ₂ O ₅)	6%
Soluble Potash (K ₂ O)	6%
Potential Acidity equivalent to 675 lbs. Calcium Carbonate per ton.	

*Equivalent to 2,4-Dichlorophenoxyacetic Acid 1.17%
**Equivalent to Silvex 0.50%

NET WT. **16** LBS.

MANUFACTURED BY

ROCKLAND CHEMICAL CO., INC.

WEST CALDWELL, N. J.



TREATS **5,000** SQ. FT.

ROCKLAND

WEED



FEED "B"

18-6-6

Fertilizes your lawn and controls weeds such as Dandelion, Buckhorn, Plantain, Burdock, Pigweed, Cocklebur, Ragweed, Chickweed and many other broadleaf lawn weeds.

Use anytime during growing season when weeds are present. Covers 5,000 sq. ft. at the following spreader settings:

SUGGESTED SPREADER SETTINGS

CYCLONE 3 SCOTT DROP 4½

JACKSON 5½ SCOTT ROTARY "B"

For smaller areas, use proportionate amounts - approximately ¾ lbs. per 1,000 square feet.

CYCLONE 3 SCOTT DROP 4½
JACKSON..... 5½ SCOTT ROTARY... "B"

For smaller areas, use proportionate amounts – approximately ¾ lbs.
per 1,000 square feet.

DIRECTIONS

Mow lawn before applying. Apply as it comes from the package. Do not mix with other materials. Do not water after application. A light dusting, broadcast by hand or spreader, is all that is necessary. If a spreader is used, wash it thoroughly before re-use.

Apply on a calm day, to avoid drifting of material as vapors from this product may injure susceptible plants in the immediate vicinity. Do not apply too close to flowers, shrubs, evergreens, etc. Be careful no application is made directly on them or over their root spread. Do not apply to St. Augustine grass, bentgrass, clover or Dichondra.

NOTE: Best results will be obtained if material is applied when grass is moist, as from morning or evening dew. Rain or watering within 24 hours after application will reduce effectiveness. Do not mow lawn until three days after treatment. Not recommended for use on newly seeded areas.

Some weeds are more resistant than others. If a repeat application is necessary, it can be made two weeks after the first application. However, lawn must be thoroughly watered before this second treatment. Areas may be re-seeded and fertilized 10 to 14 days after last application.

CAUTION: Avoid inhalation, contamination of foodstuffs and contact with skin, eyes or clothing. Wash hands after handling. Do not store near, nor allow loose material to contact seeds, fungicides, insecticides, other fertilizers. Avoid contamination of irrigation ditches or water used for domestic purposes.

This product is toxic to fish. Keep out of lakes, streams and ponds.

Do not reuse container. Destroy by burying in a safe place away from water supplies. Do not burn.