(Northo be used on Golf Courses or Sod Farms. & Statement pending litigation.

ROCKLAND

DIAZINON* SPRAY

ACTIVE INGREDIENT:

*Diazinon-Registered TM of Ciba Geigy Corp.

EPA Reg. No. 572-305

t

EPA Est. No. 572-NJ-1

ACCEPTED

JUN 3 Q 1987

15

KEEP OUT OF REACH OF CHILDREN

۰.

WARNING

See back panel for additional precautional Fungicide, and Rodenticide Act, statements. This product contains Aromatic Petroleum Solvent.

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Call a physician or Poison Control Center immediately. If possible, vomiting should be induced under medical supervision. Drink or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person. IF INHALED: Remove victim to fresh air. Apply artificial respiration if indicated. IF ON SKIN: Remove contaminated clothing. Wash affected areas with soap and water. Launder clothing before reuse. IF IN EYES: Flush eyes with plenty of water. Call a physician at once.

Net Contents 8 fl. oz., 1 p., 1 qt. Nanufactured by Rockland Chemical Co., Inc. West Caldwell, NJ 07006 (REVISED Page 2 of Label Copy 5/20/87 to be associated with submission of 5/6/87.)

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS & DOMESTIC ANIMALS: WARNING: May be fatal if swallowed. May be absorbed through the skin. Do not breathe spray mist. Avoid contact with skin, eyes and clothing. Wash thoroughly after handling and before eating or smoking. Avoid contamination of feed and foodstuffs. Do not use on household pets or humans. Do not permit children or pets to go on grass or other treated areas until spray has completely dried. Keep out of reach of domestic animals. Utensils used for measuring this product should not be reused for food preparation. PHYSICAL OR CHEMICAL HAZARDS: DO NOT USE OR STORE NEAR HEAT OR OPEN FLAME. ENVIRONMENTAL HAZARDS: This product is toxic to fish, birds, and other wildlife. Birds and other wildlife feeding in treated areas may be killed. Do not apply directly to water or wetlands. Do not apply when weather conditions favor drift om areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. 2/5

Page

2

OI

4

This product is highly toxic to bees exposed to direct treatment or +o residues remaining on treated area. Do not apply when bees are actively visiting crop, cover crop or weeds blooming in treatment area. Applications should be timed to provide maximum possible interval between treatment and next period of bee activity.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

DILUTION-SPRAYING DIRECTIONS

or

5.pt

25 ,169

آدر

₀0

.هم

įι

TOR PRESSURE OR PUMP TYPE SPRAYERS: General dilution rate 1-384. For application to Flower, Shrubs, Shade Trees, Fruits, Vegetables and Lawns: Thoroughly mix 2 tsps. of concentrate in 1 gallon of water or 2 Tbsp. (1 fl. oz.) in 3 gallons of water, or 8 fl. oz. in 24 gallons will treat 1000 sq. ft. For Millipedes in lawn areas, apply 1 pint (16 fl. oz.) in 24 gallons to 1000 sq. ft. For VEGETABLE gardens, 1 fl. oz. in 3 gallons of water will treat 1000 sq. ft.

FOR HOSE END SPRAYERS: Follow sprayer manufacturers' directions for equivalent concentrations and application rates.

FOR MELNOR SPRAY GUARD^R APPLICATION: Select setting "1" for spraying Flowers, Shrubs, Shade Trees, Fruits, Vegetables and Lawns. The 16 fl. oz. contents of this sprayer will automatically mix with water to cover 2000 sq. ft. of lawn area or 16,000 sq. ft. of Vegetables. For Millipedes, sprayer will cover 1000 sq. ft. Select Setting "2" for Outdoor Area Nuisance Insect Pests.

Rockland	Diazi.non	Spray
Page	of	

CONVERSION CHART

3 teaspoons (tsp.) = 1 Tablespoon

abel

2 Tablespoons (Tbs.) = 1 oz.

Apply when insects appear and repeat as noted in directions, or as necessary. Apply evenly to all plant surfaces. For lawns and vegetable gardens, measure area to be treated and apply at coverage rate. We suggest using a Hose End Sprayer.

LAWN SPRAY: To control Chinch bugs, Sod Webworms, Leafhoppers, Fleas, Chiggers, Milliped Ants, Brown Dog Ticks, and Earwigs. Apply a few hours after a heavy rain or thorough watering for control of all listed insects. Will also control White Grubs (Japanese Beetle larvae and European Chafer) and Billbugs. For Grubs, apply between late Jul; and early October. Water thoroughly after application. Apply for Billbugs when insects or damage first appear. For Millipedes, sprayer will cove 1,000 sq. ft. FLORAL, SHRUB AND SHADE TREE SPRAY: To control Aphids, Mites, Leafhoppers, Mealy Bugs, Thrips(Exposed stages), Leafminers and Scale Crawlers on Roses, Azaleas, Chrysanthemums, Gladioli, Carnations, Oaks, Maple, Locust, Holly, Boxwood, Pine, Spruce, Willow. Penetrate dense foliage to thoroughly cover all plant surfaces. Do not use on certain Ferns, Pointsettia, Hibiscus, Papaya, Pilea and Gardenia, since injury to the plants may occur.

FRUIT SPRAY: To control Codling Moths, Mites, Aphids, Leafrollers, and Scale Crawlers on Apples, Peaches, Plums, Grapes, Cherries, and Strawberries. Spray thoroughly to cover all plant surfaces. Start spraying after blooming and repeat at 10-14 day intervals if needed. Do not spray within 21 days of picking.

VEGETABLE SPRAY: For Tomatoes, Carrots, Fotatoes, Beans, Cucumbers, Onions, Cantelopes, Peppers and Cabbage to control Bean Beetles, Aphids, Leafminers, Imported Cabbage worms, Flea Beetles, Whiteflies, and Cucumber Beetles. Make applications at 7-10 day intervals and when insects appear. Do not spray within 10 days of picking; 35 days for Potatoes; 1 day for Tomatoes. WASH HOME GARDEN VEGETABLES BEFORE USING.

PREPLANT SOIL APPLICATIONS: Apply broadcast, just before planting at the following rates per 1000 sq. ft. in sufficient water to achieve thorough coverage. and immediately work into soil. CUTWORMS: 6 fl. oz. (Surface Cutworms, work in 2-3 in) Subterranean Cutworms, work in 3-6 in.), GARDEN SYMPHYLANS (Garden Centipedes): 15 fl. oz., work in 4-8 in., ROOT MAGGOTS (Cabbage ONLY): 45 fl. oz. work in 3-4 in., WIREWORMS: Cabbage, Carrots, Corn, Lettuce, Peas, Snap Beans and Tomatces ONLY work in 4-8 in.

3/5

Page of

> 16 32

4 i <

Dilution rates $||_{1}|_{in}$ spray window frames, doors, sills, screens, porches, patio walls, foundations, $||_{1}|_{in}$ walks, garbage cans and into cracks and other -1- μ^{μ} , walks, garbage cans and into cracks and other places where insects may hide. To μ^{μ} , control Flies. Ants Cocknowledge control Flies, Ants, Cockroaches, Boxeider Bugs, Clover Mites, Spiders, Ticks, Chiggers and Earwigs. DO NOT TREAT PLANTS with this dosage rate.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. Do not reuse empty container. Store unused material, in original container only, in a locked, dry storage area away from children and domestic animals. Wrap empty container and put in trash collection.

WARRANTY: Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

LARGER THAN ONE GALLON STORAGE AND DISPOSAL STATEMENT: Do not contaminate water, food or feed by storage or disposal. STORAGE: Store unused material, in original container only, in a locked storage area out of reach of children and animals. Keep container tightly closed. PESTICIDE DISPOSAL: Pesticide, spray mixture or rinse water that cannot be used according to label instructions must be disposed of according to applicable Federal, State or Local procedures.


CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or, puncture and dispose of in a sanitary landfill or by other procedures approved by state and local authorities.

Manufactured by:

will not

Ć

utacruted	l by:	#A30508
Rockland	Chemical Co., Inc.	16
West Cal	dwell, NJ 07006 USA	



/ บากกานบุคทบ""

3. For your protection, the top cannot be removed.