

1 2 3 4 7

ROCKLAND CO., INC.
P O BOX 503
WEST CALDWELL, N. J. 07007-0809

Front panel

ROCKLAND
ROTENONE-PYRETHRUM
INSECTICIDE

An Emulsifiable Concentrate for Insect Control on FRUITS, VEGETABLES, ORNAMENTALS
and FLOWERING PLANTS.

ACTIVE INGREDIENTS:

Rotenone.....	1.1%
Other Cube Extractives.....	2.2%
Pyrethrins.....	0.8%
Petroleum Distillate.....	3.2%
Aromatic Petroleum Solvent.....	84.7%
INERT INGREDIENTS.....	8.0%
	<hr/>
	100.0%

Contains 0.08 lb/gal. of Rotenone
Contains 0.06 lb/gal. of Pyrethrins

EPA Reg. No. 572-

EPA Est. No. 572-NJ-1

PRECAUCION AL CONSUMIDOR: Si usted no lee ingles, no use este producto hasta que la etiqueta le haya sido explicada ampliamente.
TRANSLATION: To the user: If you cannot read English, do not use this product until the label has been fully explained to you.

KEEP OUT OF REACH OF CHILDREN (12 pt. type)

WARNING (18 pt. type) AVISO

See side panel for additional precautionary statements

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Call a physician or Poison Control Center. Drink promptly a large quantity of milk, egg whites, gelatin solution, or, if these are not available, drink large quantities of water. Avoid alcohol. Do not induce vomiting. This product contains aromatic petroleum solvent. Aspiration may be a hazard.
IF IN EYES: Flush with plenty of water. Get medical attention if irritation persists.
IF INHALED: Remove from contaminated atmosphere.
IF ON SKIN: Remove contaminated clothing. Thoroughly wash effected area. Launder clothing before reuse.

NET CONTENTS:


BEST AVAILABLE COPY

ACCEPTED
WITH COMMENTS
by EPA Letter Dated:

APR 22 1985

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the
registered under

572-315



PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS: WARNING. May be fatal if swallowed. Harmful if absorbed through skin. Causes skin irritation. May cause substantial but temporary eye injury. Wear safety glasses. Do not get in eyes, on skin, or on clothing. Avoid breathing spray mist. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

ENVIRONMENTAL HAZARDS: This product is toxic to fish. Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of wastes.

PHYSICAL AND CHEMICAL HAZARDS: DO NOT USE OR STORE NEAR HEAT OR OPEN FLAME. Flash Point minimum 110°F (TCC)

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

FOR HOME GROUNDS, PARKS, RECREATIONAL AREAS, COMMERCIAL GREEN HOUSES, LATH HOUSES, and INTERIOR PLANTSCAPE to Control Insects on Shade Trees., Ornamentals and Flowering Plants. For use on such plants as poplar, linden, dogwood, willow, birch boxeider, maple, oak, elm, pine, spruce, cedar, yew, spirea, rose, lilac, zinnia, aster, petunia, geranium, dahlia, gladiolus, phlox, sweet pea, marigold, carnation, chrysanthemum, and rose. Mix this product with water as described in the following chart. Use hand or automatic spray : as a coarse spray. Use sufficient spray mix to give complete coverage of the plant. Do not apply this product within one day of collection. Repeat every 5-10 days or as needed. Keep diluted spray away from light.

USE WITH CAUTION: on African violet, azalea, Reiger begonia, browallia, Tahitian bridal veil, caenothus, camelia, plumosa fern, gloxinia, impatiens, geranium, oxalis, jade, primrose, cyclamen, wax begonia, spider plant, chrysanthemum, Boston fern, geranium, and friendship plant.

As with all plants listed above, especially carnations, chrysanthemums and roses, there exists a wide variety of species and cultivars, and therefore a wide range of susceptibility to injury. To ascertain possible phytotoxicity, spray a few plants a described below. Wait 24 hours and check for possible injury.

NOTE: Do not apply this product to plants in food service areas. Remove infested plants out of eating areas before application. They may be returned to original locations when spray is dry.

PESTS

VOLUME OF PRODUCT/GALLON OF WATER 5.79L

Aphids, Leafhoppers, Exposed Thrips	1 pt. to 100 gal. (1 tsp. to 1 gal.)
Oblique-banded Leaf Rollers, Sawfly larvae, Boxeider bugs, Japanese beetles, Elm leaf beetle larvae, Mites, Lygus bugs, Lace bugs, Greenhouse Whiteflies, Stink bugs, Garden Fleahoppers, Greenhouse Leafrollers	1 qt. to 100 gal. (2 tsp. to 1 gal.)
Cucumber beetles, Rose chafers, Ants, Aster beetles Flea beetles	2 qts. to 100 gal. (4 tsp. to 1 gal.)

BEST AVAILABLE COPY

RE-ENTRY AND ADDITIONAL APPLICATION RESTRICTIONS FOR AGRICULTURAL CROPS REQUIRING FARMWORKER HAND LABOR TASKS


Do not apply this product in such a manner as to directly, or through drift, expose workers or other persons. The area treated must be vacated by unprotected persons.

Do not enter treated areas without protective clothing until sprays have dried. Protective clothing means at least hat or other suitable head covering, a long-sleeved shirt and long-legged trousers or a coverall type garment (all closely woven fabric covering the body), shoes and socks.

Because certain states may require more restrictive re-entry intervals, for various crops treated with this product, consult your State Department of Agriculture for further information.

Written or oral warnings must be given to workers who are expected to be in treated area or in an area about to be treated with this product. Inform workers of the area or fields that must be vacated and not re-entered until the sprays have dried. Acquaint workers with the information appearing on this label under "Hazards to Humans and Domestic Animals" and "Statement of Practical Treatment." When oral warnings are given, warnings shall be given in a language best understood by workers. Oral warnings must be given if there is reason to believe that written warnings cannot be understood by workers. Written warnings must include the following information: 1) The name of this product; 2) Date of application; 3) The statement: "Do not enter without appropriate protective clothing until sprays have dried."; 4) The information given under "Hazards to Humans and Domestic Animals" and "Statement of Practical Treatment" that is appropriate in case of accidental exposure.

DIRECTIONS FOR USE ON AGRICULTURAL CROPS REQUIRING FARMWORKER HAND LABOR TASKS



BEST AVAILABLE COPY

OR use on vegetable plants such as peas, bean, best cucumber, squash, melon, potato, tomato, cabbage, spinach, lettuce, celery, asparagus, pepper, eggplant, onion, carrot and corn. Mix with water as described in the following chart. Use hand or automatic sprayer as a coarse spray. Use sufficient spray mix to give complete coverage of the plants. Do not apply this product within one day of harvest. Repeat every 5-10 days or as needed. Keep diluted spray away from light.

PESTS	VOLUME OF PRODUCT/GALLON OF WATER
Aphids, Leafhoppers, Exposed Thrips	1 pt. to 100 gal.
Cabbage Worms, Bean Beetles, Beet Webworms, Sod Webworms, Asparagus Beetles, Japanese Beetles, Lygus Bugs, Whiteflies, Garden Fleahoppers, Harlequin Bugs, Oblique-banded Leaf Rollers, Stink Bugs, Celery Leafhoppers, Melon Worms, Tortoise Beetles, Garden Webworms, Alfalfa Webworms and Mites	1 qt. to 100 gal.
Cucumber Beetles, Flea Beetles, Vegetable Weevils, Squash Bugs (nymphs), Squash Vine Borers, European Corn Borers- 3 to 4 applications or more as eggs start to hatch and repeated at 5 to 7 day intervals.	2 qts. to 100 gal.

FOR USE ON FRUITS AND BERRIES such as apple, pear, peach, plum, cherry, grape, current, gooseberry, raspberry, blackberry, strawberry and blueberry, consult the following chart. Use sufficient spray mix to give complete coverage of the plants. Do not apply this product within one day of harvest. Repeat every 5-10 days or as needed. Keep diluted spray away from light.

PESTS	VOLUME OF PRODUCT/GALLON OF WATER
Aphids, Leafhoppers, Exposed Thrips	1 pt. to 100 gal.
Imported Current Worms, Gooseberry Fruit Worms, Raspberry Fruit Worms, Blueberry Maggots, Cherry Fruit Flies, Strawberry Root Worms (adult), Red-necked Borers, Japanese beetles, and Mites	1 qt. to 100 gal.
Strawberry Leafrollers and Rose Chafers	2 qts. to 100 gal.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Store unused material, in original container only, in a locked storage area out of reach of children and animals. Keep container tightly closed.

PESTICIDE DISPOSAL: Pesticide, spray mixture or rinse water that cannot be used according to label instructions must be disposed of according to applicable Federal, State or Local procedures.

CONTAINER DISPOSAL: ONE GALLON OR LESS: Do not reuse empty container. Securely wrap in several layers of newspaper and discard in trash. Rinse thoroughly before discarding in trash. **FIVE GALLONS & LARGER -** Triple rinse (or equivalent). Then offer for recycling or reconditioning, or, puncture and dispose of in a sanitary landfill or by other procedures approved by state and local authorities.

WARRANTY: Seller makes no warranty, expressed or implied concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

BEST AVAILABLE COPY