

ACCEPTED

JUN 6 1985

Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No. 572-315

land

PYRETHRUM TICIDE

Concentrate for FRUITS, VEGETABLES, FLOWERING PLANTS.

- 1.1%
- 2.2%
- 0.8%
- 3.2%
- 84.7%
- 8.0%
- 100.00%

1/2 gal of Rotenone
3/4 of Pyrethrins

EPA EST 572 N2 1

If used no lee ingles, no ues este haya sido explicada ampliamente. Si cannot read English, do not use until fully explained to you.

PROTECTION OF CHILDREN AND THE ENVIRONMENT

ALL PRECAUTIONARY STATEMENTS

ACUTE TREATMENT

In case of poisoning, call Poison Control Center. Drink promptly a glass of water. Avoid alcohol. Do not induce vomiting. Aspirin may be harmful.

Get medical attention if irritation persists. Wash thoroughly with water after use.

NETS

PESTS	Vol. of Product Gal. of Water (3.78L)
Aphids, Leafhoppers, Exposed Thrips	1 pt. to 100 gal. — 1 tsp. to 1 gal.
Oblique banded Leaf Rollers, Sawfly larvae, Boxelder bugs, Japanese beetles, Elm leaf beetle larvae, Mites, Lygus bugs, Lace bugs, Greenhouse Whiteflies, Stink bugs, Garden Fleahoppers, Greenhouse Leafhoppers	1 qt. to 100 gal. — 2 tsp. to 1 gal.
Cucumber beetles, Rose chafers, Ants, Aster beetles, Flea beetles	2 qts. to 100 gal. — 4 tsp. to 1 gal.

ADDITIONAL APPLICATION RESTRICTIONS FOR AGRICULTURAL CROPS REQUIRING FARMWORKER HAND LABOR TASKS:

Do not apply this product in such a manner as to directly, or through drift, expose workers or other persons. The area treated must be vacated by unprotected persons.

Do not enter treated areas without protective clothing until sprays have dried. Protective clothing means at least a hat or other suitable head covering, a long-sleeved shirt and long-legged trousers or a coverall type garment (all closely woven fabric covering the body), shoes and socks.

Because certain states may require more restrictive re-entry intervals for various crops treated with this product, consult your State Department of Agriculture for further information.

Written or oral warnings must be given to workers who are expected to be in treated area or in an area about to be treated with this product. Instruct workers of the area or fields that they must be vacated and not re-entered until the sprays have dried. Acquaint workers with the information appearing on this label under "Hazards to Humans and Domestic Animals" and "Statement of Practical Treatment." When oral warnings are given, warnings shall be given in a language customarily understood by workers.

FOR USE ON VEGETABLE PLANTS such as peas, bean, beet, cucumber, squash, muskmelon, potato, tomato, cabbage, spinach, lettuce, celery, asparagus, pepper, eggplant, onion, carrot and corn. Mix with water as described in the following chart. Use hand or automatic sprayer as a coarse spray. Use sufficient spray mix to give complete coverage of the plants. Do not apply this product within one day of harvest. Repeat every 5-10 days or as needed. Keep diluted spray away from light.

PESTS	Vol. of Product Gal. of Water (3.78L)
Aphids, Leafhoppers, Exposed Thrips	1 pt. to 100 gal.
Cabbage Worms, Bean Beetles, Beet Webworms, Sod Webworms, Asparagus Beetles, Japanese Beetles, Lygus Bugs, Whiteflies, Garden Fleahoppers, Harlequin Bugs, Oblique-banded Leaf Rollers, Stink Bugs, Celery Leafhoppers, Melon Worms, Tortoise Beetles, Garden Webworms, Alfalfa Webworms and Mites	1 qt. to 100 gal.
Cucumber Beetles, Flea Beetles, Vegetable Weevils, Squash Bugs (nymphs), Squash Vine Borers, European Corn Borers - 3 to 4 applications or more as eggs start to hatch and repeated at 5 to 7 day intervals	2 qts. to 100 gal.

FOR USE ON FRUITS AND BERRIES such as apple, pear, peach, plum, cherry, grape, currant, gooseberry, raspberry, blackberry, strawberry and blueberry, consult the following chart. Use sufficient spray mix to give complete coverage of the plants. Do not apply this product within one day of harvest. Repeat every 5-10 days or as needed. Keep diluted spray away from light.

PESTS	Vol. of Product Gal. of Water (3.78L)
Aphids, Leafhoppers, Exposed Thrips	1 pt. to 100 gal.
Spotted Currant Worms, Gooseberry Fruit Worms, Raspberry Fruit Worms, Blueberry Maggots, Cherry Fruit Fly, Strawberry Root Weevil, Pear, Redstem and Borers, Japanese Beetles and Mites	1 qt. to 100 gal.
Cucumber Beetles, Flea Beetles, Vegetable Weevils, Squash Bugs (nymphs), Squash Vine Borers, European Corn Borers - 3 to 4 applications or more as eggs start to hatch and repeated at 5 to 7 day intervals	2 qts. to 100 gal.

BEST AVAILABLE COPY