

Reg # 572-315

PM-18

172

ACCEPTED
 JUL 18 1991
 572-315

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS: WARNING. May be fatal if swallowed. Harmful if absorbed through skin. Causes skin irritation. May cause substantial but temporary eye injury. Wear safety glasses. Do not get in eyes, on skin, or on clothing. Avoid breathing spray mist. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

ENVIRONMENTAL HAZARD: This pesticide is extremely toxic to fish. Do not apply directly to water or wetlands (swamps, bogs, marshes, potholes). Runoff and drift from treated areas may be hazardous to aquatic organisms in adjacent sites. Do not contaminate water when disposing of equipment washwaters.

PHYSICAL AND CHEMICAL HAZARD: DO NOT USE OR STORE NEAR HEAT OR OPEN FLAME. Flash Point minimum 110°F.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Store unused material, in original container only, in a locked storage area out of reach of children and animals. Keep container tightly closed.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: ONE GALLON OR LESS: Do not reuse empty container. Securely wrap in several layers of newspaper and discard in trash. Rinse thoroughly before discarding in trash. **FIVE GALLONS & LARGER** - Triple rinse (or equivalent). Then offer for recycling or reconditioning, or, puncture and dispose of in a sanitary landfill or by other procedures approved by state and local authorities.

WARRANTY: Seller makes no warranty, expressed or implied concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

FOR HOME GROUNDS, PARKS, RECREATIONAL AREAS, COMMERCIAL GREEN HOUSES, LATH HOUSES, and INTERIOR PLANTSCAPE to Control insects on Shade Trees, Ornamentals and Flowering Plants. For use on such plants as poplar, linden, dogwood, willow, birch, boxelder, maple, oak, elm, pine, spruce, cedar, yew, spirea, rose, lilac, zinnia, aster, petunia, geranium, dahlia, gladiolus, phlox, sweet pea, marigold, carnation, chrysanthemum, and rose. Mix this product with water as described in the following chart. Use hand or automatic sprayer as a coarse spray. Use sufficient

Rockland.

ROTENONE-PYRETHRUM INSECTICIDE

An Emulsifiable Concentrate for Insect Control on
FRUITS, VEGETABLES, ORNAMENTALS and FLOWERING PLANTS.

ACTIVE INGREDIENTS:

Rotenone.....	1.1%
Other Cube Extractives.....	2.2%
Pyrethrins.....	0.8%

*INERT INGREDIENTS.....	95.9%
	100.00%

*Contains xylene range aromatic solvent.

Contains 0.06 lb/gal of Rotenone; 0.06 lb/gal of Pyrethrins

EPA Reg. No. 572-315

EPA Est. 572-NJ-1

PRECAUCION AL CONSUMIDOR: Si usted no lee ingles, no use este producto hasta que la etiqueta le haya sido explicada completamente.

TRANSLATION: (To the user: If you cannot read English, do not use this product until the label has been fully explained to you.)

KEEP OUT OF REACH OF CHILDREN WARNING

SEE BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Call a physician or Poison Control Center. Drink promptly a large quantity of milk, egg whites, gelatin solution, or, if these are not available, drink large quantities of water. Avoid alcohol. Do not induce vomiting. This product contains aromatic petroleum solvent. Aspiration may be a hazard.

IF IN EYES: Flush with plenty of water. Get medical attention if irritation persists.

IF INHALED: Remove from contaminated atmosphere.

IF ON SKIN: Remove contaminated clothing. Thoroughly wash affected area. Launder clothing before reuse.

NET CONTENTS:

292

spray mix to give complete coverage of the plant. Do not apply this product within one day of collection. Repeat every 5-10 days or as needed. Keep diluted spray away from light.

Use with caution in greenhouses and on interior plants: African violet, azalea, Reiger begonia, bromelia, Tahitian bridal veil, caenothus, camellia, plumosa fern, gloxinia, impatiens, geranium, oxalis, jade, primrose, cyclamen, wax begonia, spider plant, chrysanthemum, Boston fern, geranium, and friendship plant.

As with all plants listed above, especially carnations, chrysanthemums and roses, there exists a wide variety of species and cultivars, and therefore a wide range of susceptibility to injury. To ascertain possible phytotoxicity, spray a few plants as described below. Wait 24 hours and check for possible injury.

NOTE: Do not apply this product to plants in food service areas. Remove infested plants out of eating areas before application. They may be returned to original locations when spray is dry.

GENERAL ARBORICULTURAL APPLICATION: Use 1-4 pts. per 100 gals. for controls of leaf feeding homoptera, coleoptera and lepidoptera insects.

MIST BLOWER APPLICATION: Follow Spray Equipment Manufacturer's directions for equivalent dilution rates.

PESTS

Aphids, Leafhoppers, Exposed Thrips

Vol. of Product
Gal. of Water (3.78L)

1 pt. to 100 gal. —
1 tsp. to 1 gal.
1 qt. to 100 gal. —
2 tsp. to 1 gal.

Oblique-banded Leaf Rollers, Sawfly larvae, Boxelder bugs, Japanese beetles, Elm leaf beetle larvae, Mites, Lygus bugs, Lace bugs, Greenhouse Whiteflies, Stink bugs, Garden Fleahoppers, Greenhouse Leafhoppers

Cucumber beetles, Rose chafers, Ants, Aster beetles, Flea beetles

2 qts. to 100 gal. —
4 tsp. to 1 gal.

ADDITIONAL APPLICATION RESTRICTIONS FOR AGRICULTURAL CROPS REQUIRING FARMWORKER HAND LABOR TASKS:

Do not apply this product in such a manner as to directly, or through drift, expose workers or other persons. The area treated must be vacated by unprotected persons.

Do not enter treated areas without protective clothing until sprays have dried. Protective clothing means at least a hat or other suitable head covering, a long-sleeved shirt and long-legged trousers; or a coverall type garment (all closely woven fabric covering the body), shoes and socks. Because certain states may require more restrictive re-entry intervals for various crops treated with this product, consult your State Department of Agriculture for further information.

Written or oral warnings must be given to workers who are expected to be in treated area or in an area about to be treated with this product. Instruct workers of the area or fields that they must be vacated and not re-entered until the sprays have dried. Acquaint workers with the information appearing on this label under "Hazards to Humans and Domestic Animals" and "Statement of Practical Treatment." When oral warnings are given, warnings shall be given in a language customarily understood by workers.

FOR USE ON VEGETABLE PLANTS such as peas, bean, beet cucumber, squash, muskmelon, potato, tomato, cabbage, spinach, lettuce, celery, asparagus, pepper, eggplant, onion, carrot and corn: Mix with water as described in the following chart. Use hand or automatic sprayer as a coarse spray. Use sufficient spray mix to give complete coverage of the plants. Do not apply this product within one day of harvest. Repeat every 5-10 days or as needed. Keep diluted spray away from light.

PESTS

Aphids, Leafhoppers, Exposed Thrips
Cabbage Worms, Bean Beetles, Beet Webworms, Sod Webworms, Asparagus Beetles, Japanese Beetles, Lygus Bugs, Whiteflies, Garden Fleahoppers, Marquis Bugs, Oblique-banded Leaf Rollers, Stink Bugs, Celery Leafhoppers, Melon Worms, Tortoise Beetles, Garden Webworms, Alfalfa Webworms and Mites

Vol. of Product
Gal. of Water (3.78L)

1 pt. to 100 gal.
1 qt. to 100 gal.

Cucumber Beetles, Flea Beetles, Vegetable Weevils, Squash Bugs (nymphs), Squash Vine Borers, European Corn Borers — 3 to 4 applications or more as eggs start to hatch and repeated at 6 to 7 day intervals.

2 qts. to 100 gal.

FOR USE ON FRUITS AND BERRIES such as apple, pear, peach, plum, cherry, grape, current, gooseberry, raspberry, blackberry, strawberry and blueberry, consult the following chart. Use sufficient spray mix to give complete coverage of the plants. Do not apply this product within one day of harvest. Repeat every 5-10 days or as needed. Keep diluted spray away from light.

PESTS

Aphids, Leafhoppers, Exposed Thrips
Imported Current Worms, Gooseberry Fruit Worms, Raspberry Fruit Worms, Blueberry Maggots, Cherry Fruit Flies, Strawberry Root Worms (adult), Red-necked Borers, Japanese beetles, and Mites
Strawberry Leafrollers and Rose Chafers

Vol. of Product
Gal. of Water (3.78L)

1 pt. to 100 gal.
1 qt. to 100 gal.

2 qts. to 100 gal.

Manufactured by
ROCKLAND CORPORATION
West Caldwell, N.J. 07006

B315

BEST AVAILABLE COPY