

Reg no: 572-316

PM-17

132

Rockland. MILL-MIST 'S'



A synthetic Pyrethroid for controlling the following household flying and crawling insects: Houseflies, Mosquitoes, Gnats, Fruit Flies, Wasps, Hornets, Yellow Jackets, Cockroaches, Bedbugs, Fleas, Ants, Crickets, Silverfish, Earwigs, Spiders, Grain Insects (adults of Confused flour beetles, Saw-toothed grain beetles, Shiny spider beetles), Grain mites, Sowbugs, Centipedes and Millipedes

ACTIVE INGREDIENTS

* Resmethrin [5 (phenylmethyl) 3 (uranyl) methyl 2,2 dimethyl 3 (2 methyl 1 propenyl) cyclopropanecarboxylate	0.250%	w/w
Related compounds	0.034%	
Petroleum distillate	99.706%	
INERT INGREDIENTS	0.010%	

Authorized by USDA for use in Federally inspected meat, poultry and egg processing plants.

* Cis/trans isomers ratio max 30% (±) cis and min 70% (±) trans

* US Patent Nos. 3,465,007 and 3,542,928

EPA Reg. No. 572.316

EPA Est. 572 NJ 1

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

See Side Panel For Additional Precautionary Statements

NET CONTENTS:

STATEMENT OF PRACTICAL TREATMENT
IF SWALLOWED Call a physician or Poison Control Center. Do not induce vomiting. Drink large quantities of milk, egg whites or gelatin solution. If these materials are not available, drink large quantities of water. This product contains petroleum solvents. Aspiration may be a hazard.
IF IN EYES Flush with plenty of water.
IF ON SKIN Remove contaminated clothing. Wash affected area with soap and water. Launder clothes before re-using.
IF INHALED Remove victim to fresh air.

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION Harmful if swallowed or absorbed through skin. Avoid contact with eyes, skin, or clothing. Avoid breathing mist. Cover or remove exposed foods. Avoid contamination of cooking utensils and food preparation surfaces. Remove pets and cover fish aquariums before spraying.
ENVIRONMENTAL HAZARDS Do not spray on turf or plants.
PHYSICAL OR CHEMICAL HAZARDS Extinguish open flame before space spraying. This formulation is combustible. **DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME.**

STORAGE AND DISPOSAL
STORAGE Store unused material, in original, closed container only, in a locked, dry storage area out of reach of children and animals. Keep container tightly closed. Do not contaminate water, food or feed by storage or disposal. Do not use, pour, spill or store near heat or open flame.
PESTICIDE DISPOSAL Pesticide wastes are toxic. Improper disposal of excess pesticide or pesticide rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.
CONTAINER DISPOSAL ONE GALLON — Do not reuse empty container. Securely wrap in several layers of news paper and discard in trash. Rinse thoroughly before discarding in trash. **FIVE GALLONS & LARGER** — Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill or by other procedures approved by state and local authorities.
WARRANTY Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling that is contrary to label instructions.

ACCEPTED
 1977 17 1980
 572-316

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling

FOR USE INDOORS AND OUTSIDE PREMISES OF HOTELS, MOTELS, HOMES, FOOD HANDLING ESTABLISHMENTS, PACKAGING AND PROCESSING PLANTS, SEED HOUSES, COMMERCIAL, INSTITUTIONAL AND STORAGE BUILDINGS, MILK HANDLING AREAS OF MILK PLANTS, MILK STORAGE ROOMS, EGG PROCESSING PLANTS, SUPERMARKETS, RESTAURANTS, BOTTLING PLANTS, CANNERIES, FLOUR AND FEED MILLS, BAKERIES, BOATS, SHIP CABINS AND HOSPITALS IN HOSPITALS. PATIENTS SHOULD BE REMOVED PRIOR TO TREATMENT Also for use in bushes, automobiles, trucks, boxcars and ship cargo holds. Also for use in dog kennels and horse stables. In Horse Stables, not for application if horses will be used for food

Food should be removed or covered during treatment. All food processing surfaces should be covered during treatment or thoroughly cleaned before using. When using the product in these areas apply only when the facility is not in operation

INDOORS: SPACE SPRAY To kill houseflies, mosquitos, and gnats. Close room or area to be treated. Shut off air conditioner and ventilator system. Seal all open vents. Remove fish and pets. Apply undiluted as a very fine mist space spray through manually operated equipment or with electrical and other powered applicators. Avoid staining by holding the spray nozzle at least 5 ft from objects. Direct the fine spray mist toward the upper corners of the room. Fill the room with a fine mist space spray at the rate of 5 fl ozs per 5,000 cu ft. Leave room closed for one-half hour. Ventilate thoroughly before re entry. Sweep up and destroy insects

CONTACT SPRAY To kill cockroaches, fleas, ants, crickets, silverfish, earwigs, spiders, grain mites, sowbugs, centipedes and millipedes. Adjust spray to produce large wetting type spray particles. Direct spray to contact these insects in cracks and crevices in floors, baseboards, around pipes, behind cabinets, and other harborages. Spray these infested areas until thoroughly wet. Repeat as necessary

TO KILL BEDBUGS Apply as a spot treatment to cracks and crevices around baseboards, floorboards, bed

boards and walls. First, thoroughly clean and air mattresses and springs. Then spot treat only the areas that may harbour bedbugs, concentrating on tufts, folds and seams. Repeat as necessary. For persistent infestations, consult professional applicator

TO KILL FLEAS Hold spray nozzle about 18 inches from areas to be treated. Direct spray on resting areas and floor around these areas. Apply spray until area is moist. Treat pet bedding the same way. Also spray entire inside surface of dog houses. Repeat as necessary. Do not allow pets to contact treated areas until spray has dried. Remove or cover animal feed before spraying. Also, treat dogs, cats and horses with a registered flea control product before they re-enter treated area

TO KILL WASPS, HORNETS, YELLOW JACKETS Application should be made in late evening when insects are at rest. Spray liberally into hiding and breeding places contacting as many insects as possible. Repeat as necessary

TO KILL SPIDERS Spray directly on these pests and use repeated applications to the webbing and all cracks and crevices which harbor spiders

TO KILL ANTS Spray doors, around window frames, ant hills and trails

TO KILL GRAIN INSECTS Adults of confused flour beetles, sawtoothed grain beetles shiny spider beetles. Remove infested grain, cereal or flour and destroy. Avoid contamination of other food and foodstuffs in the areas by removing before spraying. Spray directly on exposed stages and into cracks and crevices where grain insects may be hiding. In homes, dispose of all infested products. The shelves and drawers should be cleaned and thoroughly sprayed so as to contact all pests, and covered with shelf paper after allowing shelves to dry. Any utensils that are contacted with spray should be washed before using. Repeat as necessary

THERMAL APPLICATIONS This formulation is to be used undiluted in only such portable thermal applicator equipment that can deliver spray particle sizes within the aerosol size range

For the control of houseflies, mosquitoes and gnats in warehouses, restaurants and industrial institutions. Remove or cover exposed food handling surfaces. Remove animals and plant life. Shut off ventilating systems, air ducts, air conditioning equipment and close off room or area to be treated. Do not apply while food processing plant is occupied in operation

Extinguish all flames and pilot lights. Apply as a dry fog at the rate of 1-1/3 fl ozs per 1,000 cu ft of space. Direct nozzle of thermal applicator toward the upper corners of the room and along the ceiling. Close off room for at least one half hour. Do not remain in treated areas. Ventilate the area after treatment. Sweep up and destroy fallen insects. Repeat treatment as necessary

OUTDOORS: To control nuisance pests around buildings, patios and paved areas. Apply as a contact spray. Spray directly on insects. Avoid spraying turf or plants

ANTS AND FLEAS Same as directed for indoor use
WASPS, HORNETS, YELLOW JACKETS Same as directed for indoor use. Aim spray at nest openings in bushes, under eaves, chimneys, cracks and crevices to saturation in and around nest openings. Contact as many insects as possible

EARWIGS, SOWBUGS, SPIDERS, CENTIPEDES AND MILLIPEDES Apply as a contact spray, as for indoor use. To control named flying and crawling insects around premises spray thoroughly around doors, window frames and similar areas of entry. Also thoroughly spray garbage and refuse areas especially around poultry houses, barns and similar areas of heavy infestation. Do not spray on turf or plants
THERMAL APPLICATIONS This formulation is to be used undiluted in only such portable thermal applicator equipment that can deliver spray particle size within the aerosol size range

For reduction in annoyance from houseflies, mosquitoes, and gnats in yards, patios, around swimming pools, picnic areas, camp sites, drive ins and horse stables, and other **SMALL AREA** applications

Hold thermal applicator at least 5 ft away from any object, especially plants and shrubbery. Avoid wetting any plant foliage. Apply undiluted at the rate of 3 1/2 gallons per acre. Fog only when there is no breeze, or the wind velocity is less than 3 m p h. The ideal time is generally at dusk when the wind dies down and the air temperature and ground temperature are approximately the same. Under these conditions the fog will linger in the treated area for a longer time. Apply fog while moving away from treated area. Avoid inhalation of the fog. Do not enter the treated area until the fog has dissipated. Repeat application as necessary