		\$ 1.50 P. 1.50	•		• •	- 326	PJ 167	
Please read instructions on reverse before completing form. United States Environmental Protecti Washington, DC 20		F A	ome No. 207 Registration Comendme Other	n	OPP Identifier Number 229235			
Application for Pesticide - Section I								
1. Company/Product Number 572-326		2. EPA Product Menager 3. Proposed Classification				, !		
4. Company/Product (Name) ROCKLAND Rockland Home Pest Spray (Alt) FLY-RID		PM# X None Restricted						
Name and Address of Applicant (Include ZIP Cade) ROCKLAND CORPORATION P.O. Box 809 West Caldwell, NJ 07006 Check if this is a new address		6. Expedited Review. In accordance with FIFRA Section 3(c)(3) (b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. Product Name						
Section - II								
Amendment - Explain below. Resubmission in response to Agency letter dated X Notification - Explain below. Explanation: Use additional page(s) if necessary. (For section of the context of the contex	Final printed labels in response to Agency letter dated "Me Too" Application. Other - Explain below.							
Update effluent discharge labeling statement per PR 95-1 required on packaging of 5 gallons and larger.								
·	Sect	ion - III						
1. Meterial This Product Will Be Packaged In:	- 1 -							
Child-Resistant Packaging Yes No * Certification must be submitted Unit Packaging Yes No No No Packaging Ves No No No Packaging Ves No No Packaging Ves No No Packaging Ves No No Packaging Ves No No Packaging		No Pl GI			Metal Plastic Glass Paper	etal astic ass		
Label Container		tail Container 5. Location of .abel Directions On Label On Labeling accompanying product					uct	
S. Manner in Which Label is Affixed to Product Lithograph Paper glued Stenciled Other								
Section - IV								
1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.),								
Judith K. Lemm	Mg	r., Regulatory Affairs		fairs	Telephone No. (Includ's Area Code) 201-575-1322			
Certification I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law. 6. Date Application Received (Stamped) (Stamped)								
2. Signature Hemm		Mgr., Regulatory Affairs						
4. Typed Name Judith K. Lemm	5, Date	6/30/95						

ROCKLAND

FLY-RID

(Alt Names: Home Pest Spray, Watch-Em-Drop)

A concentrated PERMENONE(R) Stable Emulsion CONTACT and SPACE SPRAY.

- For Control of Insects Effecting HORSE, DAIRY and BEEF CATTLE, GOATS, SHEEP & HOGS.
- For Use outdoors and in HOMES and NON-FOOD AREAS, COMMERCIAL BUILDINGS, WAREHOUSES, THEATERS, OFFICE BUILDINGS, SCHOOLS, MOTELS, HOTELS and FOOD/FEED PROCESSING ESTABLISHMENTS and KENNELS.
 (Not for Use in U.S. Department of Agriculture (USDA) Meat & Poultry Plants)

ACTIVE INGREDIENT:

*Permethrin	0.5%
INERT INGREDIENTS	99.5%
	100.00%

^{*(3-}phenoxyphenyl) methyl (+/-) cis/trans 3-(2,2-dichloroethenyl) 2,2-dimethyl-cyclopropanecarboxylate. cis/trans ratio: Min. 35% (+/- and max. 65% (+/-)trans.

(R)PERMANONE is a Reg. Trademark of Roussel Uclaf Corp.

EPA Reg. No.: 572-326

EPA Est. No. 572-NJ-1

KEEP OUT OF REACH OF CHILDREN

CAUTION

See Side Panel for Additional Precautionary Statements

Net Contents:

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS: Harmful if swallowed. May be absorbed through skin. Avoid inhalation of spray mist. Avoid contact with skin, eyes, or clothing. Wash thoroughly after handling and before smoking or eating. Avoid contamination of feed or food stuffs. Remove pets and cover fish tanks before spraying. Do not use on household pets or humans. Do not allow children or pets to contact treated areas until surfaces are dry.

ENVIRONMENTAL HAZARDS: This product is toxic to fish and aquatic organisms. For terrestrial uses, do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water by disposing of equipment washwaters: When using 5 gallon container or larger for commercial, institutional or non-food areas of Food Processing establishments, do not discharge effluent containing this product into lakes, streams, ponds, estuaries, oceans or other waters unless in accordance with the requirements of a National Pcludant Discharge Elimination System (NPDES) permit and the permitting authority has been notified in writing prior to discharge. Do not discharge effluent containing this product to sewer systems without previously notifying the local sewage treatment plant authority. For guidance contact your State Water Board or Regional Office of the EPA>

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Drink one or two glasses of water and induce vomiting by touching the back of the throat with finger. Repeat until vomit fluid is clear. Call a physician immediately. Do not induce vomiting or give anything by mouth to an unconscious person.

IF INHALED: Remove affected person to fresh air. Apply artificial respiration if indicated

IF ON SKIN: Remove contaminated clothing and wash affected areas with soap and water.

IF IN EYES: Flush eyes with plenty of water. Call a physician if irritation persists.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Not for use on plants being grown for sale or other commercial use, or for commercial seed production, or for research purposes. For use on plants intended for aesthetic purposes or climatic modification and being grown in interior plantscapes, ornamental gardens or parks, or on golf courses or lawns and ground.

SHAKE WELL BEFORE USING.

LIVESTOCK AND FARM USE

To protect Horses, Cattle (beef & dairy), Goats, Sheep, and Hogs, from Horn Flies, Face Flies, House Flies, Mosquitoes and Gnats: Apply a light mist sufficient to wet the surface of the hair. For face flies, spray face and head, but do not spray into eyes.

To Control Stable Flies, Horse Flies and Deer Flies: Apply at a rate of 2 ounces per adult animal, sufficient to wet the hair thoroughly. Repeat treatment daily or at intervals necessary to give continued protection.

To Control Blood-Sucking Lice: Apply to the infested areas of the animal, using a stiff brush to get the spray to the base of the hair. Repeat every 3 weeks if required.

To Control Poultry Lice: Spray roosts, walls and nests or cages thoroughly. This should be followed by spraying over the birds with a fine mist.

To Control Bedbugs and Mites in Poultry Houses: Spray crevices of roost poles, cracks in walls, and cracks in nests where the bedbugs and mites hide.

ON DOGS: To control Fleas, Ticks and Lice: Start spraying at the tail, moving the dispenser rapidly and making sure that the animal's entire body is covered, including the legs and underbody. While spraying, fluff the hair so that the spray will not penetrate to the skin. Make sure spray wets thoroughly. Do not spray into eyes or face. Avoid contact with genitalia. Repeat as necessary. NOTE: Do not apply on puppies less than four weeks old.

PREMISE TREATMENTS: BARNS, MILK HOUSES, OUT BUILDINGS, PENS Space and Surface applications should be made separately from direct animal applications.
Animals should be removed from treated area. Do not contaminate feed, drinking water,
milk, or milking equipment. Vacate treated areas. Ventilate before reoccupying.
SPACE SPRAY: To control Stable flies, Horse flies, House flies, Deer flies, Mosquitoes,
Gnats, Angoumois Grain Moths, Mediterranean Flour Moths and Indian Meal Moths.
Close doors and windows and shut off ventilating systems. Adjust application equipment
to deliver a fine mist or fog. Direct spray at an upward angle to obtain uniform
distribution throughout the entire area at a rate of 1 fl. Oz. per 1000 cubic ft. Keep area
closed for two hours before ventilating.

SURFACE SPRAY: To control Cockroaches, Spiders, Flies, Gnats, Exposed stages of Grain Moths. Apply as a coarse, wet spray to walls, ceilings, around windows and to other surfaces where insects congregate. Treat cracks and crevices where insects hide. For Ants, spray trails, nests and points of entry. Contact as many insects as possible. For Surface Treatment ONLY, use at the rate of 1 gallon per 750 sq. ft.

FOR INDOOR APPLICATIONS IN HOMES, INSTITUTIONS AND RESTAURANTS

The use of this product in food processing establishments should be confined to time periods when the plant is not in operation. Food should be removed or covered during

treatment. All food processing surfaces should be covered during treatment and washed with an effective cleaning compound followed by a potable water rinse before using.

Vacate treated areas and ventilate before reoccupying. For maximum effectiveness, a combination of spot, surface treatment and fogging is recommended.

SURFACE SPRAYING: To Control cockroaches, waterbugs, palmetto bugs, scorpions, millipedes, centipedes, sowbugs, pillbugs, ants, silverfish, firebrats, spiders, crickets, clover mites, cheese mites, granary weevils, rice weevils, confused flour beetles, rust, red flour beetles, drugstore beetles, meal worms, grain mites and cadelles: Use a good sprayer adjusted to a coarse wet spray. Direct the spray into hiding places, cracks and crevices, under pallets, around containers of stored foods, around the base of machinery, and behind shelves and drawers. For silverfish, spray bookcases. For ants, spray trails, nests and points of entry. If surface application only is to be used, spray floors and other surfaces applying at a rate of 1 gallon to 750 square feet of surface.

To Control Carpet Beetles: Spray edges of carpeting and under carpeting and rugs. Make localized application to floor and baseboards. Spray directly into cracks, crevices and infested areas of shelving. Repeat treatment as needed.

To Control Bedbugs: Spray mattresses lightly, particularly around tufts and seams. Take beds apart and spray in all joints. Treat baseboards, moldings and floors. Repeat treatment as needed.

To control Fleas and Ticks (adults and larvae): Thoroughly spray infested areas, pet beds, resting quarters, nearby cracks and crevices, along and behind baseboards, moldings, window and door frames, and entire areas of floor and floor covering. Fresh bedding should be placed in animal quarters following treatment. Repeat treatment as needed.

To Control Clothes Moths: Remove any infested articles from storage, brush thoroughly, and air for several hours in sunlight, if possible. Apply spray liberally to empty chests, closets, bureaus and other storages directing the spray into cracks, joints and crevices. After airing, infested articles may be treated lightly, keeping nozzle at least three feet from fabric to avoid staining.

To Kill Lice and Louse Eggs: Spray in an inconspicuous area to test for possible staining or discoloration. Inspect again after drying, then proceed to spray entire area to be treated. Spray from a distance of 8 to 10 inches. Treat only those garments and parts of bedding, including mattresses and furniture that cannot be either laundered or dry cleaned. Allow all treated articles to dry thoroughly before use.

SPACE SPRAYING: Close doors and windows and shut off ventilating systems. Apply with a good machine adjusted to deliver a fine mist or fog. For rapid control of house flies, fruit flies, gnats, mosquitoes, skipper flies, wasps, hornets, bees, black flies, Angoumois grain moths, Mediterranean flour moths, Indian meal moths and tobacco

moths: Direct spray at an upward angle, distributing it uniformly throughout the entire area at a rate of 1 ounce per 1000 cubic feet of space. Keep area closed for 2 hours. Ventilate before reoccupying.

FOGGING: To kill exposed or accessible stages of the above named crawling insects, use a rate of 1 gallon to 20,000 to 40,000 cubic feet of space. Contact insects directly when possible. Keep area closed for 2 hours. Ventilate before reoccupying.

OUTDOOR APPLICATION

FOR CONTROL OF GARDEN INSECTS: For use on roses, dahlias, asters and other ornamentals to kill Japanese beetles hit by spray as well as whiteflies, aphids, exposed thrips, spiders, red mites and leafminers.

May be used on plants as a multi-purpose spray in small gardens and for spot treatments of incipient insect infestations to prevent spreading in large garden areas.

May also be used on other plants such as African violets, asters, azaleas, begonias, camellias, carnations, chrysanthemums, geraniums, crassula, grape vines, Kentia palms, laurels, marigolds, rhododendrons, roses, rubber plants, snapdragons, stocks, wandering Jews and zinnias.

Effective when used as directed against whiteflies, aphids, spider and red mites, exposed thrips, lace bugs, Japanese beetles, pavement ants, clover mites and army worms.

When spraying plants, do not operate closer than 18 inches. Use sweeping motion. Be cautious about wetting tender foliage, young plants and new growth. Do not spray plants when temperatures exceed 90 F. Contact insects directly whenever possible. Repeat treatment as infestations accord.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Store in a cool, dry area away from heat or open flame. Store above 32° F. PESTICIDE DISPOSAL: Pesticide, spray mixture, or rinse water that cannot be used according to label instructions must be disposed of at or by an approved waste disposal facility.

CONTAINER DISPOSAL: LARGER THAN 1 GALLON: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke. Wrap container in several layers of newspaper and discard in the trash.

WARRANTY: Seller makes no warranty, expressed or implied concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

Manufactured by:
Rockland Corporation
West Caldwell, NJ

Product C32601,05
0594 (wps exc.
statement added)
6/94 Mosquitoes
deleted-per DCI(unsupported)