

9/16/97

195

SEP 16 1997

~~3354-52797~~
174

Ms. Judith K. Lemm
Rockland Corporation
P.O. Box 809
West Caldwell, NJ 07006

Dear Ms. Lemm:

Subject: Amendment- delete use on dogs/product name change
Rockland Indoor/Outdoor Insect Sprāy
EPA Registration No. 572-350
Your Submission dated July 30, 1997

This amendment referred to above, submitted in connection with registration under FIFRA section 3(c)(7)(a), is acceptable, provided that you:

1. Submit and/or cite all data or other material required for registration/reregistration of your product under FIFRA section 3(c)(5) or FIFRA section 4 when the Agency requires all registrants of similar products to submit such data.

2. Make the labeling changes listed below before you release the product for shipment bearing the amended labeling:

a. Under Directions for Use add "Do Not Use on Dogs or Cats".

3. Submit two (2) copies of your final printed labeling before you release the product for shipment. Refer to the A-79 enclosure for a further description of final printed labeling.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

Sincerely,

George T. LaRocca
Product Manager (13)
Insecticide-Rodenticide Branch
Registration Division (7505C)

Enclosure

New, Proposed LABEL. 295

Rockland Indoor/Outdoor Insect Spray

- For Domestic Use Both Indoors and Outdoors.
- A Ready-To-Use PERMANONE® Stable Emulsion Insecticide.
- For Control Of Fleas, Brown Dog Ticks, and Lice On Premises.
- For Control of Pests on Livestock.
- Provides 60-Day (Two Months) Control of Cockroaches, Fleas and Stored Product Pests.

ACTIVE INGREDIENT:

*Permethrin.....0.25%

INERT INGREDIENTS.....99.75%
100.00%

*(3-phenoxyphenyl) methyl (+/-) cis/trans 3-(2,2-dichloroethenyl) 2,2-dimethylcyclopropanecarboxylate. *Cis/trans* ratio: min.35%(+/-)*cis* and max. 65% (+/-) *trans*.

PERMANONE -Registered Trademark of AgrEvo Environmental Health, Inc.

KEEP OUT OF REACH OF CHILDREN

CAUTION

See Side Panel for Additional Precautions

Net Contents: One Quart (32 Ounces), One Gallon

ACCEPTED
with COMMENTS
In EPA Letter Dated

SEP 10 1997

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

572-350

READ ENTIRE LABEL BEFORE EACH USE

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS. CAUTION: Wash thoroughly after handling and before smoking or eating. Avoid contamination of feed and foodstuffs. Remove pets and birds and cover fish aquaria before surface applications. Do not use on humans. Do not allow children or pets to contact treated areas until surfaces are dry.

ENVIRONMENTAL HAZARDS: This product is extremely toxic to fish. Do not apply directly to any body of water. Do not contaminate water when disposing of equipment washwaters.

FIRST AID: If Swallowed, do not induce vomiting. Call physician or Poison Control Center immediately. If In Eyes, flush with plenty of water. Get medical attention. If On Skin, remove contaminated clothing. Wash affected area with soap and water. Launder clothing before reuse. If Inhaled, remove from contaminated atmosphere.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. Do not reuse container. Wrap container in several layers of newspaper and discard in trash.

Affix trigger, pump or other type sprayer to container for application. Do not transfer to another another container.

SHAKE WELL BEFORE USING.

INDOOR APPLICATION

For spot application only. Do not use as a space spray indoors. In the home, cover all food handling surfaces and cover or remove all food and cooking utensils, or wash thoroughly after treatment.

Surface Spraying: For 60 day (2 months) control of Cockroaches, Waterbugs, Palmetto Bugs, Ants, Silverfish, Firebrats, Spiders, Crickets, Clover Mites, Cheese Mites, Granary Weevils, Rice Weevils, Confused Flour Beetles, Rust Red Flour Beetles, Drugstore Beetles, Meal Worms, Grain Mites and Cadelles: Direct the spray into hiding places, cracks and crevices and behind shelves and drawers. For Silverfish, spray bookcases. For Ants, Spray trails, nests and points of entry.

To Control Carpet Beetles: Spray edges of carpeting and under carpeting and rugs. Make localized application to floor and baseboards. Spray directly into cracks, crevices and infested areas of shelving. Repeat treatment as needed.

To Control Bedbugs: Spray mattresses lightly, particularly around tufts and seams. Take beds apart and spray in all joints. Treat baseboards, moldings and floors. Repeat treatment as needed.

To Control Fleas (Adult and Larvae) and Ticks (Adult and Larvae): Thoroughly spray infested areas, pet beds, resting quarters, nearby cracks and crevices, along and behind baseboards, moldings, window and door frames, and entire areas of floor and floor covering. Fresh bedding should be placed in animal quarters following treatment. Repeat treatment as needed. Allow to dry thoroughly before allowing pets to contact treated areas.

Do Not Use on Cats or Dogs

To Control Clothes Moths: Remove any infested articles from storage, brush thoroughly, and air for several hours in sunlight, if possible. Apply spray liberally to empty chests, closets, bureaus and other storages directing the spray into cracks, joints, and crevices. After airing, infested articles may be treated lightly, keeping nozzle at least three feet from fabric to avoid staining.

To Kill Lice and Louse Eggs: Spray in an inconspicuous area to test for possible staining or discoloration. Inspect again after drying, then proceed to spray entire area to be treated. Spray from a distance of 8 to 10 inches. Treat only those garments and parts of bedding, including mattresses and furniture that cannot be either laundered or dry cleaned. Allow all treated articles to dry thoroughly before use.

ON LIVESTOCK: To protect cattle (beef and dairy), goats, sheep, hogs and horses from

Horn Flies, Face Flies, House Flies, Mosquitoes and Gnats: Apply a light mist sufficient to set the surface of the hair. For face flies, spray face and head, but do not spray into eyes.

To Control Stable Flies, Horse Flies and Deer Flies: Apply at a rate of 2 ounces per adult animal, sufficient to wet the hair thoroughly. Repeat treatment daily or at intervals necessary to give continued protection.

To Control Blood-Sucking Lice: Apply to the infested areas of the animal, using a stiff brush to get the spray to the base of the hair. Repeat every 3 weeks if requires.

To Control Poultry Lice: Spray roosts, walls and nests or cages thoroughly. This should be followed by spraying over the birds with a fine mist.

To Control Bedbugs and Mites in Poultry Houses: Spray roosts, walls and nests or cages thoroughly. This should be followed by spraying over the birds with a fine mist.

5 9 5
4

OUTDOOR APPLICATION

To help control ticks in outdoor activity areas, treat a band of vegetation and leaf litter 6-10 feet wide and adjacent to activity areas, especially where dense vegetation occurs.

For Control of Garden Insects: For use on roses, dahlias, asters and other ornamentals to kill Japanese Beetles hit by spray as well as Whiteflies, Aphids, Exposed Thrips, Spiders, Red Mites and Leafminers. May be used on plants as a multi-purpose spray in small gardens and for spot treatments of incipient insect infestations to prevent spreading in large garden areas.

May also be used on other plants such as African violets, asters, azaleas, begonias, camellias, carnations, chrysanthemums, dahlias, delphiniums, dogwoods, English ivy, eunonymus, fuschia, geraniums, crassula, grape vines, Kentia palms, laurels, marigolds, rhododendrons, roses, rubber plants, snapdragons, stocks, wandering jews and zinnias.

Effective when used as directed against Whiteflies, Aphids, Spider and Red Mites, exposed Thrips, Lace Bugs, Japanese Beetles, Pavement Ants, Clover Mites and Armyworms. When spraying plants, do not operate closer than 28 inches. Use sweeping motion. Be cautious about wetting tender foliage, young plants and new growth. Do not spray plants when temperatures exceed 90 F. Contact insects directly whenever possible. Repeat treatment as infestations occur.

WARRANTY: Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

Manufactured by:
Rockland Corporation
West Caldwell, NJ 07006
Product A35032

EPA Reg. No. 572-350
EPA Est. No. 572-NJ-1