

PM 22

572-354

7/30/96

17 1/10



U.S. ENVIRONMENTAL PROTECTION AGENCY
Office of Pesticide Programs
Registration Division (7505C)
401 "M" St., S.W.
Washington, D.C. 20460

EPA Reg. Number:
572-354

Date of Issuance:
JUL 30 1996

Term of Issuance:
Conditional

Name of Pesticide Product:
Rockland Ornamental Fungicide

NOTICE OF PESTICIDE:
 X Registration
 Reregistration
(under FIFRA, as amended)

Name and Address of Registrant (include ZIP Code):

Rockland Corporation
686 Passaic Avenue
P.O. Box 809
West Caldwell, NJ 07007-0809

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA sec. 3(c)(7)(A) provided that you:

1. Submit and/or cite all data required for registration/reregistration of your product under FIFRA sec. 3(c)(5) when the Agency requires all registrants of similar products to submit such data; and submit acceptable responses required for reregistration of your product under FIFRA section 4.

2. Make the following label changes listed below before you release the product for shipment:

a. Add the phrase, "EPA Reg. No. 577-354".

b. Since this product is in Toxicity Category III, CAUTION, the Spanish Language Text and English Translation are not required to appear on the labeling of this product. Also, the statement "Stop - Read the Label Before Use" is not required by our regulations and can be deleted from the labeling.

Signature of Approving Official:

ASJ

Date:

JUL 30 1996

page 2
EPA Reg. No. 577-354

c. The statements "Not for use on plants being grown for sale....or lawns and grounds" are not required to appear on the front panel but should be placed in close proximity to "For Control of Certain Diseases on Flowers, Lawns, Shrubs, Shade Trees and Turf". This section should be visible to a consumer when purchasing the product.

3. Submit one (1) copy of your final printed labeling before you release the product for shipment. Refer to the A-79 enclosure for a further description of final printed labeling.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

Cynthia Giles-Parker
Product Manager (22)
Fungicide-Herbicide Branch
Registration Division (7505C)

Enclosure

3/10

ACCEPTED
with COMMENTS
In EPA Letter Dated

JUL 30 1996

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No. 572-354

(Front Panel of Jar Label)

Rockland Ornamental Fungicide

with Bayleton®



For Control of Certain Diseases on Flowers, Lawns, Shrubs, Shade Trees, and Turf.
Not for use on plants being grown for sale or other commercial use, or for commercial seed product, or for research purposes. For use on plants intended for aesthetic purpose or climatic modification and being grown in interior plantscapes, ornamental gardens, yards or home gardens or on golf courses or lawns and grounds.

ACTIVE INGREDIENT:	
1-(4-Chlorophenoxy)-3, 3-Dimethyl-1-(1H-1,2,4- triazol-1-yl)-2-butanone.....	25%
INERT INGREDIENTS.....	75%
	<hr/>
	100%

STOP-READ THE LABEL BEFORE USE
KEEP OUT OF REACH OF CHILDREN
CAUTION
See Rear Panel for complete Precautionary Statements

PRECAUCION AL USUARIO: Si usted no puede leer o entender ingles, no use este producto hasta que la etiqueta le haya sido explicada ampliamente. (TO THE USER: If you cannot read or understand English, do not use this product until the label has been fully explained to you.)

U.S. Patent No. 3,912,752

Net Weight 2 Ounces

®Bayleton is a registered Trademark of Bayer Ag

(Back Panel of Jar Label:)

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS: CAUTION: Harmful if swallowed, inhaled, or absorbed through the skin. Causes eye irritation. Prolonged or frequently repeated skin contact may cause allergic skin reactions in some individuals. Avoid breathing dust. Avoid contact with eyes, skin, or clothing. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse

STATEMENTS OF PRACTICAL TREATMENT: IF SWALLOWED: Call a physician or Poison Control Center. Vomiting should be induced. Administer water freely and induce vomiting by giving one dose (1/2 oz. or 15 ml.) of syrup of ipecac. If vomiting does not occur within 10 to 20 minutes, administer second dose. If syrup of ipecac is not available, drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Repeat until vomit fluid is clear. Do not induce vomiting or give anything by mouth to an unconscious person., Professional medical assistance should be secured immediately.

IF IN EYES; Flush with plenty of water. Call a physician if irritation persists. **IF ON SKIN:** Wash with plenty of soap and water. Get medical attention. **SYMPTOMS OF POISONING:** The compound does not cause any definite symptoms that would be

4/10
b

diagnostic. Poisoning is accompanied by hyperactivity followed by sedation. **NOTE TO PHYSICIAN:** No specific antidote. Treat symptomatically.


ENVIRONMENTAL HAZARDS: Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters. Apply this product only as specified on this label. This chemical demonstrates the properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Do not make applications when weather conditions favor drift from target area.

FOR DIRECTIONS FOR USE, STORAGE AND DISPOSAL , READ SUPPLEMENTAL LABELING ACCOMPANYING THIS CONTAINER.

EPA Reg. No. 572-_____
Est. No. 572-NJ-1
Product A36502 0196

Manufactured by:
Rockland Corporation
West Caldwell, NJ 07006 USA

UPC



(Copy for Hang-Tag on 2 oz. jar.)

Rockland Ornamental Fungicide with Bayleton®

For Control of Certain diseases on Flowers, Lawns, Shrubs, Shade Trees and Turf..

ACTIVE INGREDIENT:

1-(4-Chlorophenoxy)-3,3-dimethyl-1-H-1,2,4-triazol-1-yl)-2-butanone.....	25%
INERT INGREDIENTS.....	75%
	100%

STOP-READ THE LABEL BEFORE USE

KEEP OUT OF REACH OF CHILDREN
CAUTION

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS:CAUTION: Harmful if swallowed, inhaled, or absorbed through the skin. Causes eye irritation. Avoid breathing dust. Avoid contact with eyes, skin, or clothing. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.


STATEMENTS OF PRACTICAL TREATMENT:

IF SWALLOWED: Call a physician or Poison Control Center. Vomiting should be induced. Administer water freely and induce vomiting by giving one dose (1/2 oz. or 15 ml.) of syrup of ipecac. If vomiting does not occur within 10 to 20 minutes, administer second dose. If syrup of ipecac is not available, drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Repeat until vomit fluid is clear. Do not induce vomiting or give anything by mouth to an unconscious person., Professional medical assistance should be secured immediately. **IF IN EYES:** Flush with plenty of water. Call a physician if irritation persists. **IF ON SKIN:** Wash with plenty of soap and water. Get medical attention. **SYMPTOMS OF POISONING:** The compound does not cause any definite symptoms that would be diagnostic. Poisoning is accompanied by hyperactivity followed by sedation. **NOTE TO PHYSICIAN:** No specific antidote. Treat symptomatically.

ENVIRONMENTAL HAZARDS: Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters. Apply this product only as specified on this label. This chemical demonstrates the properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Do not make applications when weather conditions favor drift from target area.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.



IMPORTANT: Read these entire Directions and Conditions of Sale before using this product.

Not for use on plants being grown for sale or other commercial use, or for commercial seed production, or for research purposes. For use on plants intended for aesthetic purposes or climatic modification and being grown in interior plantscapes, ornamental gardens, parks or home gardens or on golf courses or lawns and grounds.

CONDITIONS OF SALE: THE DIRECTIONS ON THIS LABEL WERE DETERMINED THROUGH RESEARCH TO BE APPROPRIATE FOR THE CORRECT USE OF THIS PRODUCT. THIS PRODUCT HAS BEEN TESTED UNDER DIFFERENT ENVIRONMENTAL CONDITIONS BOTH INDOORS AND OUTDOORS UNDER CONDITIONS SIMILAR TO THOSE THAT ARE ORDINARY AND CUSTOMARY WHERE THE PRODUCT IS TO BE USED. INSUFFICIENT CONTROL OF PESTS OR PLANT INJURY MAY RESULT FROM THE OCCURRENCE OF EXTRAORDINARY OR UNUSUAL CONDITIONS, OR FROM FAILURE TO FOLLOW LABEL DIRECTIONS. IN ADDITION, FAILURE TO FOLLOW LABEL DIRECTIONS MAY CAUSE INJURY TO ANIMALS, MAN, AND DAMAGE TO THE ENVIRONMENT. ROCKLAND OFFERS, AND THE BUYER ACCEPTS AND USES, THIS PRODUCT SUBJECT TO THE CONDITIONS THAT EXTRAORDINARY OR UNUSUAL ENVIRONMENTAL CONDITIONS, OR FAILURE TO FOLLOW LABEL DIRECTIONS ARE BEYOND THE CONTROL OF ROCKLAND AND ARE, THEREFORE, THE RESPONSIBILITY OF THE BUYER.

APPLICATION: Rockland Ornamental Fungicide is absorbed rapidly and works systemically from within the plant. Good coverage and wetting of the foliage are necessary. Rainfall or sprinkler irrigation within 30 min. after application does not decrease effectiveness. Control may be less effective on plants suffering from drought stress. Therefore, in order to achieve maximum control, plants should be maintained in a vigorously growing state through good cultural practices. Apply in all cases when plants are fully established and actively growing. Applications should be applied at recommended intervals to maintain disease control. DO NOT USE ON CROPS GROWN FOR FOOD OR FORAGE. DO NOT APPLY THIS PRODUCT THROUGH ANY TYPE OF IRRIGATION SYSTEM.

See Charts For Application Rates And Diseases Controlled.

COMPATIBILITY: Rockland Ornamental Fungicide is compatible with many registered insecticides and fungicides. To determine the compatibility of ornamental fungicides this product with specific products, the following procedure should be conducted. Pour the recommended proportions of the products into a suitable container of water, mix thoroughly and allow to stand at least 5 minutes. If the combination remains mixed or can be remixed readily, the mixture is considered physically compatible.

SPRAY ADDITIVES: Use of various spray additives such as spreaders, extenders, trace elements or fertilizers should be evaluated prior to use. The label directions given here are based on data obtained with no additives; use of any product with Ornamental Fungicide may affect the result. Contact local university extension personnel prior to use of spray mix additives.

RESTRICTIONS: Edible portions of treated trees, other than apples and pears, such as nuts and syrup, should not be used for feed or food.

STORAGE AND DISPOSAL: Do not contaminate water, food or feed by storage or disposal. Store in a cool dry place and in such a manner as to prevent cross contamination with other pesticides, fertilizers, food, and feed. Store in original container and out of the reach of children, preferably in a locked storage area.

Handle and open container in a manner as to prevent spillage. If the container is leaking or material spilled for any reason or cause, carefully sweep material into a pile. Refer to Precautionary Statements on label for hazards associated with the handling of this material. Do not walk through spilled material. Material that cannot be used as directed should be disposed of as directed below. In spill or leak incidents, keep unauthorized people away. You may contact Chemtrec at 800-424-9300 for decontamination procedures or any other assistance that may be necessary.

HOUSEHOLD: Do not reuse empty container. Securely wrap in original container in several layers of newspaper and put in trash collection.

WARRANTY: Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions

Manufactured by : EPA Reg. No. 572-XXX
Rockland Corporation EPA Est. No. 572-NJ-1
West Caldwell, NJ 07006 Product A 36502 1095

"BAYLETON" is a Registered Trademark of Bayer Ag

ORNAMENTAL PLANT DISEASE CONTROL

Locate plant(s) (see below) to be treated. Cross reference the number/letter codes, following the plant name, to the specific diseases (see below) controlled. Refer to Application Rates section for instructions detailing use for each disease. In California, only those plants marked with an asterisk may be treated.

NOTE CONVERSION: 7.08 TABLESPOONS OR 21.24 TEASPOONS = ONE OUNCE

PLANTS	Diseases
Flowering & Foliage Plants (Outdoor)	(1) Flower Blight
Ageratum (2b,3,4)	(a) <i>Ovulinia</i> spp. [A]
Aster (4)	(b) <i>Sclerotinia</i> spp. [A]
Begonia* (3)	(c) <i>Oidium</i> spp. [A]
Canna (4)	
Carnation (3,4)	
Chrysanthemum (3,4)	(2) Leaf Blight/Spots
Dahlia (3)	(a) <i>Cephalosporium</i> spp. [C]
Delphinium (3)	(b) <i>Cercospora</i> spp.
Dianthus (4)	(c) <i>Didymella</i> spp. [B]
Four O'Clock (4)	(d) <i>Didymascella thujina</i> [G]
Geranium* (3,4)	(e) <i>Entomosporium</i> spp. [C]
Hollyhock* (3,4)	(f) <i>Exobasidium</i> spp. [E]
Hydrangea (3)	
Iris* (2c)	(3) Powdery Mildew
Marigold (2b,4)	<i>Erysiphe</i> spp.
Nephthytis* (2a)	<i>Microsphaera</i> spp.
Pansy (3,4)	<i>Oidium</i> spp.
Petunia (3,4)	<i>Podosphaera</i> spp.
Phlox (2b,3,4)	<i>Phyllactinia</i> spp.
Poinsettia (3)	<i>Sphaerotheca</i> spp.
Rose* (3)	<i>Uncinula</i> spp.
Salvia (3,4)	(4) Rusts
Sedum (3)	(a) <i>Coleosporium</i> spp.
Snapdragon* (3,4)	(b) <i>Cronartium</i> spp. [B]
Sunflowers(3,4)(ornamental only)	(Fusiform)
Sweet Peas* (3)	(c) <i>Gymnosporangium</i> spp.
Zinnia* (2b,3)	(d) <i>Melampsora</i> spp. [F]
	(e) <i>Melampsora farlowii</i> [A]
	(f) <i>Melampsoridium</i> spp.
	(g) <i>Peridermium</i> spp. [B]
	(h) <i>Phragmidium andersonii</i>
	(i) <i>Puccinia</i> spp.
	(j) <i>Uromyces</i> spp.
	(k) <i>Uredinopsis mirabilis</i> [A]
	(5) Tip Blight
	<i>Sirococcus strobilinus</i> [B]
SHADE TREES	
Ash (3)	
Aspen (3,4)	
Birch (3,4)	
Buckeye (3)	
Chestnut (3)	
Cottonwood (3,4)	
Elm (3)	
Fir (4)	
Locust (3)	
Maple (3)	
Oak* (3)	
Pine* (4,5)	
Poplar (3,4)	
Russian Olive 2b,4)	
Sycamore* (3)	
Walnut (3)	
Willow* (3,4)	
Ornamental Shrubs & Trees	
Amelanchier (3)	
Azalea* (3)	
Barberry (3,4)	
Buckthorn (4)	
Camellia (Suppres. of 1b)	
Cedar* (2d)	
Crabapple (flowering) (3,4)	
Crape myrtle* (3)	
Dogwood (3)	
Euonymus* (3)	
Gardenia (3)	
Hawthorn (3,4)	
Hemlock (4e)	
Holly (3)	
Juniper (4)	
Leucothoe (2b)	
Lilac (3)	
Mock-Orange (3,4)	
Mountain Laurel (1a,2b,3)	
Ninebark (3)	
Paulownia (3)(Empress Tree)	
Pear (flowering) (3)	
Photinia (2e,3,4)	
Potentilla (4) (Cinquefoil)	
Privet (2b,3)	
Pyracantha (3)	
Rhododendron (1a,2b,3)	
Spirea (3)	
Viburnum* (3,4)	
Vitex (2b) (Chaste Tree)	
Flowering & Foliage Plants	
Non-commercial Greenhouse (D)	
African Violet* (3)	
Azalea (1a,2f,3)	
Calendula (3,4)	
Carnation* (3,4)	
Chrysanthemum* (3,4)	
Cineraria (3)	
Crassula (3)	
Daisy (3,4)	
Fern, Boston (4) Desmella spp.	
Geranium* (3,4)	
Gerbera (3)	
Grape Leaf Ivy* (3)	
Hydrangea (3)	
Kalanchoe (3)	
Poinsettia (3)	
Rose* (3)	
Snapdragon (3,4)	

Application Rates: Except as noted for specific diseases, mix 1/2 tsp. to 1 tsp of Ornamental Fungicide per gallon of water (1 to 2 ounces in 50 gallons of water) and apply as a full coverage foliage spray to the point of drip as needed.

[A] Mix 2.8 tsp. to 5.67 tsp. of Rockland Ornamental Fungicide in 5 gals. of water (4 to 8 ounces in 50 gallons of water) and apply as a full coverage foliar spray to the point of drip. Applications should begin at the expanded bud stage (color showing). Use multiple applications at 7 to 14 day intervals as needed dependent upon bloom periods.

[B] Mix 5.67 tsp. Rockland Ornamental Fungicide plus sufficient spreader sticker for good coverage in 5 gallons of water(8 ounces in 50 gallons of water). Apply in a spray application to the point of run-off on an as needed basis during the early part of the season. Excessive rates or excessive applications may result in a shortening of the flower stalk on iris.

[C] Mix 2.8 tsp to 5.67 tsp of Rockland Ornamental Fungicide in 5 gallons of water (4 to 8 ounces in 50 gallons of water) and apply as a full coverage foliar spray to point of run-off. Apply in early spring as growth starts and re-apply on a 14 to 21 day interval until new growth is fully expanded. Protect new growth that develops in late summer or fall as temperatures begin to drop

[D] Non-commercial Greenhouse Applications: Winter use- .8 tsp of Ornamental Fungicide in 5 gallons of water (1 oz in 50 gallons of water). Summer Use- 1.6 tsp. of Rockland Ornamental Fungicide in 5 gallons of water (2 ounces in 50 gallons of water).

[E] For control of *Exobasidium* flower and leaf gall, apply 1.6 tsp. of Rockland Ornamental Fungicide in 5 gallons of water (2 ounces in 50 gallons of water). Begin application at bud break and apply at 10 day intervals through infestation period.

[F] For control of *Melampsora pinitorqua* (Pine Twisting Rust), apply a single application in spring during periods favorable for infection. Mix 5.67 tsp. Rockland Ornamental Fungicide in 5 gallons of water (16 ounces in 50 gallons of water) and apply to shoots in the upper whorl of suspectable pine species. Make a single application per year as a full coverage application sprayed to runoff.

[G] For control of *Didymascella thujina*, Cedar Leaf Blight, apply 5.67 tsp. Rockland Ornamental Fungicide per 10 gallons of water) 0.5 lb.per 100 gallons) applied as a full coverage spray to ornamentals. Begin applications before disease appears in spring, and repeat at 60 day intervals through early fall.

NOTE CONVERSION: 7.08 TABLESPOONS OR 21.24 TEASPOONS = ONE OUNCE

RECOMMENDED APPLICATION FOR TURF GRASS DISEASE CONTROL

DISEASE	OZ. OF ROCKLAND ORNAMENTAL FUNGICIDE PER 1,000 SQ. FEET*	REMARKS
Anthraxnose (<i>Colletotrichum graminicola</i>)	Preventive 2	PREVENTIVE RATE: Apply at 30 day intervals and repeat as necessary for seasonal control. Dependent upon environmental conditions, residual control may be extended to 45 days. PREVENTIVE RATE: (Except California) Apply at 21-day intervals and repeat as necessary for seasonal control. Dependent upon environmental conditions, residual control may be extended to 30 days. CURATIVE RATE: To control existing infections, apply the curative rate. Subsequent applications should be applied on a preventive schedule and rate.
	1	
	--	
Southern Blight (<i>sclerotium rolfsii</i>)	Preventive 1 to 4	PREVENTIVE: Begin applications prior to the appearance of disease symptoms. Depending on anticipated disease severity, apply 2 to 4 oz. rates at 14 day intervals for the initial 2 to 3 treatments. Apply subsequent treatments of 1 to 2 oz. at 14 to 28 day intervals. CURATIVE: To control existing infections, apply 4 oz. at 14 day intervals for the initial 2 to 3 treatments followed by 1 to 2 oz. at 14 to 28 day intervals.
	Curative 4	
Stripe Smut (<i>Ustilage striiformis</i>)	Preventive Rates Only 2	Make a single application in spring as grass growth begins.
Fusarium Blight (<i>Fusarium culmorum</i>) (<i>Fusarium poae</i>)	2 to 4	Apply first application in the Spring 30 to 60 days before initial symptoms normally appear. Repeat applications at 30 day intervals as needed.
Summer Patch (<i>Magnaporthe poae</i>)	2 to 4	Make first application in early fall (mid Sept. to mid Oct.) prior to development of disease symptoms. A second application in early spring may be necessary in areas where disease pressure is known to be heavy.
Zoysia patch Large patch of zoysia (<i>Rhizactonia solani</i>)	2 to 4	Make first application in early fall (mid Sept. to mid Oct.) prior to development of disease symptoms. A second application in early spring may be necessary in areas where disease pressure is known to be heavy.
Dollar Spot (<i>Sclerotinia homoeocarpa</i>)	Preventive 1	PREVENTIVE RATE: Apply recommended rate on 30 day intervals. Protective activity of this product may extend for as long as 60 days dependent upon environmental conditions. PREVENTIVE RATE: (Except California) Apply recommended rate on 14 day intervals. Protective activity of this product may extend for as long as 30 days dependent upon environmental conditions. CURATIVE RATE: To control existing infections, apply the curative rate. Subsequent applications should be applied on a preventive schedule and rate.
	0.5	
	--	
Copper Spot (<i>Gloeocercospora sorghi</i>)	1	PREVENTIVE RATE: Apply at 15 to 30 day intervals. When environmental conditions favor light to moderate diseased development, use a longer interval. Protective activity of this product can be greater than 30 days depending on environmental conditions.
Powdery Mildew (<i>Erysiphe graminis</i>)	1	CURATIVE RATE: To control existing infections, apply the curative rate. Subsequent applications should be applied on a preventive schedule and rate.
Corticium Red Thread (<i>Laetisaria fuciformis</i>)	1	PREVENTIVE RATE: Apply at 15 to 30 day intervals. When environmental conditions favor light to moderate diseased development, use a longer interval. Protective activity of this product can be greater than 30 days depending on environmental conditions.
Rusts (<i>Puccinia spp.</i>)	1	CURATIVE RATE: To control existing infections, apply the curative rate. Subsequent applications should be applied on a preventive schedule and rate.
Brown Patch/ Rhizoctonia Blight (<i>Rhizoctonia solani</i>)	1	PREVENTIVE RATE: Apply at 15 to 30 day intervals. When environmental conditions favor light to moderate diseased development, use a longer interval. Protective activity of this product can be greater than 30 days depending on environmental conditions.
Gray Snow Mold/Typhula Blight (<i>Typhula incarnate</i>)	Preventive Rates Only 4	Apply in the fall, 30 days prior to turf dormancy. If turf breaks dormancy during winter months, a second cover, or when turf is dormant.
Pink Snow Mold/Fusarium Patch (<i>Microdochium nivalis</i>)	2 to 4	Apply before conditions favorable for infection occur. Re-application should be made as needed at a 60 to 90 day interval. Do not apply over snowcover, or when turf is dormant. Use higher rate in areas with a history of severe disease damage.

NOTE: *Apply the specified amount of ornamental fungicide in 2 to 4 gallons of spray per 1,000 sq. ft. for equivalent rates per acre see conversion table below. Make all applications after mowing and allow foliage to dry thoroughly before irrigation. Do not use clippings for animal feed.

RECOMMENDED APPLICATIONS (CONTINUED)
TURF GRASS DISEASE CONTROL

DISEASE	OF ROCKLAND ORNAMENTAL FUNGICIDE PER 1,000 SQ. FEET*		REMARKS
Bermudagrass decline (<i>Gaeumannomyces graminis</i> var. <i>graminis</i>)	Preventive 2	Curative 4	Immediately after the fungicide is applied, the area should be thoroughly irrigated to move the active ingredient down into the crown and root zone of the turf. The amount of water is dependent on the depth of the root zone. The objective is to water the fungicide into the crown and root zone. PREVENTIVE RATE: Begin applications 2 to 4 weeks prior to the appearance of disease symptoms. Initiate cultural control practices at the same time the fungicide is applied. Refer to your local County Extension Service for this information. Apply subsequent applications at 21 to 28 day intervals. CURATIVE RATE: To control existing infections, apply 4 oz at 14 to 21 day intervals for the initial 1 or 2 treatments followed by 2 oz. at 21 to 28 day intervals. Cultural control practices such as aeration, topdressing, reseeding, and fertilization must be implemented prior to or at the same time the fungicide is applied. Refer to your local County Extension Service for this information.

*NOTE: Apply the specified amount of this product in 2 to 4 gallons of spray per 2,000 sq. ft. For equivalent rates per acre see Conversion Table below. Make all applications after mowing and allow foliage to dry thoroughly before irrigation. Do not use clippings for animal feed.

Conversion Table for Rockland Ornamental Fungicide

OZ. per 1000 sq. ft.	==	Lb per Acre
0.5	==	1.4
1	==	2.75
2	==	5.5
4	==	11.5
5	==	13.75

Amount of BAYLETON 25% Wettable Powder to Use in Volume of Water				
Gallons of Water				
50	25	10	5	1
.5 oz	7 gm	2.3 gm	1.4 gm	.3 gm
1 oz	.5 oz	5.7 gm	2.3 gm	.8 gm
2 oz	1 oz	11.3 gm	5.3 gm	1.1 gm
3 oz	1.5 oz	17.0 gm	8.4 gm	1.7 gm
4 oz	2 oz	.3 oz	11.2 gm	2.3 gm
8 oz	4 oz	1.3 oz	.8 oz	4.5 gm
1/4 level teaspoonful equals 0.3 grams 1/2 level teaspoonful equals 0.5 grams 1 level teaspoonful equals 1.3 grams 1 level tablespoonful equals 4 grams 3 level teaspoonfuls equals 1 level tablespoonful 28.35 grams equals 1 oz				