

577-97

01/29/48

1/2

Acme Bordeaux Mixture

A convenient Dry Bordeaux Mixture, of well-balanced formula, uniform in composition, having excellent suspension and adhesive qualities.

THIS PACKAGE CONTAINS:

Active Ingredient: Copper, expressed as metallic, (in the form of Bordeaux Mixture).....	12.75%
Inert Ingredients.....	87.25%
	<hr/> 100.00%



POISON



Antidote for Copper: Give large quantities of white of eggs or milk. Call a doctor.

Seller makes no warranty of any kind, express or implied, concerning the use of this product. Buyer assumes all risk of use or handling whether in accordance with directions or not.

ACME WHITE LEAD AND COLOR WORKS

*Administration Offices: Detroit, Michigan
Factories and Warehouses in Principal Cities*

Sold for Fungicide Purposes



POISON



UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTER-
ED UNDER NO. 577-97

Jan. 29, 1948

DIRECTIONS

For the control of Blotch, Bitter Rot, Black Rot, and Anthracnose of Apples; Brown Rot of Apricots, Plums (except Japanese varieties), and Prunes; Black Rot of Grapes; and Blight, Flea Beetle, and Leaf Hopper of Potatoes, use at standard strength.

For the control of Peach Leaf Curl apply Acme Bordeaux Mixture at the rate of 12 to 16 tablespoonfuls per gallon of water. Application must be made in the Fall or early Spring when the tree is dormant.

For Strawberry Leaf Spot, Tomato Leaf Spot, Celery Blight, Brown Rot of Lemons, Scab of Grapefruit, and Melanose of Oranges and Grapefruit, use at low strength.

For control of certain Leaf Spots and Blights of Delphinium, Geranium, Laurel, Lilac, and Rhododendron, use at $\frac{3}{4}$ of the low strength on the chart. For Hollyhock, Ivy, Lily, Phlox, and Leaf Blotch of Peony, use at low strength. For Boxwood, Chrysanthemum, and Iris Leaf Spots, use at standard strength.

NOTE.—Not to be used for spraying such tender foliage as that of Peaches and Japanese Plums, nor for early growing period spraying of Apples. When used on Apples after the early growing period against the diseases specified, add hydrated lime at the rate of four pounds per fifty gallons of diluted spray.

Mixing Table

Use level spoonfuls in measuring, 3 teaspoonfuls equal 1 tablespoonful.

Amount of Spray Desired	Amount of Bordeaux Mixture Required	
	Low Strength	Std. Strength
1 Pint	$\frac{3}{4}$ to 1 tablespoon	1 to $1\frac{1}{2}$ tablespoons
1 Quart	$1\frac{1}{4}$ to $1\frac{3}{4}$ tablespoons	2 to 3 tablespoons
1 Gallon	6 to 7 tablespoons	8 to 12 tablespoons
8 Gallons	$1\frac{1}{4}$ to $1\frac{3}{4}$ cupfuls	$1\frac{1}{2}$ to $2\frac{1}{4}$ cupfuls

For more complete directions—
write for Acme Spraying Guide
Made in U.S.A.