



POISON

Acme Bordeaux Mixture

Acme's Dry Bordeaux Mixture, a balanced formula, uniform in suspension, having excellent suspension and adhesive qualities.

THIS PACKAGE CONTAINS:

12.5% of Copper, expressed as available (the form of Bordeaux Mixture) 57.5% net weight.

POISON
Antidote for Copper: Give large quantities of white of eggs or milk. Call a doctor.

Seller makes no warranty of any kind, expressed or implied, concerning the use of this product. Buyer assumes all risk in handling whether in accordance with directions or not.

WHITE LEAD AND COLOR WORKS

Manufacturers of White Lead, Detroit, Michigan
Sole U.S. Distributors in Philadelphia, Pennsylvania

MARCH
MAR 2, 1949
UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTERED
UNDER NO. 577-97

DIRECTIONS

For the control of Black, Bitter Rot, Black Rot, and Anthracnose of Apples; Brown Rot of Apples, Plums (except Japanese varieties), and Prunes; Black Rot of Grapes; and Blight, Flea Beetle, and Leaf Hopper of Potatoes, use at standard strength.

For the control of Peach Leaf Curl apply Acme Bordeaux Mixture at the rate of 12 to 16 tablespoons per gallon of water. Application must be made in the fall or early Spring when the tree is dormant.

For Strawberry Leaf Spot, Tomato Leaf Spot, Calary Blight, Brown Rot of Lemons, Scab of Grapefruit, and Melanose of Oranges and Grapefruit, use at low strength.

For control of certain Leaf Spots and Blights of Delphinium, Geranium, Laurel, Lilac, and Rhododendron, use at 1/4 of the low strength on the leaves. For Hollyhock, Ivy, Lily, Peony and Leaf Blotch of Peony, use at low strength. For Boxwood, Chrysanthemum, and Iva Leaf Spots, use at standard strength.

NOTE—Not to be used for spraying such tender foliage as that of Peonies and Japanese Plums, nor for early spring spraying of Apples. When used on Apples after the early growing period, use the standard strength, and hydrated lime at the rate of four pounds per 100 gallons of diluted spray.

Mixing Table

Use level spoonfuls in measuring. 8 teaspoonfuls equal 1 tablespoonful.

Amount of Spray Desired	Amount of Bordeaux Mixture Required	
	Low Strength	Std. Strength
1 Pint	1/4 to 1/2 tablespoon	1 to 1 1/2 tablespoons
1 Quart	1/2 to 3/4 tablespoon	2 to 3 tablespoons
1 Gallon	1 1/4 to 2 1/4 cups	3 to 6 cups

WRITE FOR LATEST SPRAYING AND DUSTING GUIDE

MACEY U.S.A.