


577-97

08/16/60

1/6

(Front panel)



Active Ingredient: Copper,
 expressed as metallic,
 (in the form of Bordeaux
 Mixture.....12.75%
 Inert Ingredients.....87.25%
100.00%

Made for
 Acme Quality Paints, Inc.
 Detroit 11, Michigan

Made in U.S.A.

LD119-06 8-59 2279

ONE
FOUND

NET
WEIGHT

(Revised
5 July, 60)



(Right Side panel)
ACME BORDEAUX MIXTURE

DIRECTIONS
(8-8-100 is the same as 4-4-50)

For the control of Blotch, Bitter Rot, Black Rot and Late Scab of Apples; Brown Rot of Apricots, Plums (except Japanese varieties) and Prunes; Black Rot and Blight of Grapes; early and late Blight of Potatoes; and the repelling of Flea Beetles and Leafhoppers on Potatoes. Use 8-8-100. Apply in spring. Repeat as necessary to maintain control. See Note.

For Pecan Scab, use at 4-4-100 when catkins show. Repeat 3 or 4 times at 3-week intervals.

For control of Peach Leaf Curl. Apply fall or early spring when tree is dormant. Use 8-8-100 at 12 tablespoonfuls per gallon. See Note.

For Strawberry Leaf Spot, Tomato Leaf Spot, Celery Blight, Brown Rot of Lemons, Scab of Grapefruit and Melanose of Oranges and Grapefruit, use at 4-4-100.


For control of Spanish Moss, spray during winter period. Use at 8-8-100.

For die back on Camellias and Botrytis Blight (gray mold) on Lilies, Dahlias, Tulips and Gladiolus, use at 4-4-100 in early spring.

MIXING TABLE

Use level spoonfuls in measuring. 3 teaspoonfuls equal 1 tablespoonful. Use higher dosage if can has been shaken to fluff the powder.

Amount of Spray Desired	Amount of Bordeaux Mixture Required	
	4-4-100	8-8-100
1 Pint	3/4 to 1 tablespoon	1 to 1-1/2 tablespoons
1 Quart	1-1/2 to 1-3/4 tablespoons	2 to 3 tablespoons
1 Gallon	6 to 7 tablespoons	8 to 12 tablespoons
3 Gallons	1-1/8 to 1-1/4 cupfuls	1-1/2 to 2-1/4 cupfuls



(Back panel)
ACME BORDEAUX MIXTURE

For control of certain Leaf Spots and Blights of Delphinium, Geranium, Laurel, Lilac and Rhododendron, use at 4 table-spoonfuls per gallon.

For Leaf Spot, Scab and Anthracnose on Palms; Anthracnose and Leaf Spot of Hollyhock; Leaf Spot of Ivy; Leaf Blotch of Peony; use at 4-4-100.

For Iris Leaf Spot, Dogwood Leaf Spot, Boxwood Nectria Canker and Chrysanthemum Leaf Spot, use 8-8-100 as growth starts in spring. Repeat if necessary.

For Needle Cast on Blue Spruce and Sphaeropsis Blight of Pine, use 8-8-100.

For Coryneum Blight of Cypress, Juniper and Arborvitae, apply fall and spring at 4-4-100.


For Twig Blight of Yew, apply in spring as growth starts. *on Cedars use in July-August*

For Anthracnose, Leaf Spot and Elm Leaf Curl, *for earlier apply* spray Sycamores, Linden, Maple, Oak and Elms *rest.* as leaves uncurl in spring. Repeat in 2 to 3 weeks.

WRITE FOR LATEST SPRAYING

AND DUSTING GUIDE

Sold For Fungicide Purposes



(Left panel)

ACME BORDEAUX MIXTURE

Dry Bordeaux Mixture is made from equal parts of Copper Sulphate and Lime. Eight pounds of this product, mixed with 50 gallons of water, is equivalent to a 4-4-50 Bordeaux formula. This strength is sometimes referred to as 8-8-100.

Disease Control is a matter of prevention. Application should be made before hand or just as soon as first signs of disease appear.

A convenient Dry Bordeaux Mixture, of well-balanced formula, uniform in composition, having excellent suspension and adhesive qualities.

NOTE--Not to be used for spraying such tender foliage as that of Peaches and Japanese Plums, nor for early growing period spraying of Apples. When used on Apples after the early growing period against the diseases specified, add hydrated lime at the rate of four pounds per fifty gallons of diluted spray.

NOTICE:--Neither the manufacturer nor the seller makes any representation or warranty, express or implied, and shall not be held responsible in any manner for any personal injury, property damage or other type of loss resulting from the handling, storage or use of this product, whether or not used in accordance with directions.

CAUTION: HARMFUL IF SWALLOWED.



ACME BORDEAUX MIXTURE

MIXING TABLE
Level spoonfuls in measuring, 3 tea-
equal 1 tablespoonful.

Amount of Bordeaux Mixture Required	
Low Strength	Std. Strength
1/4 to 1 tablespoon	1 to 1 1/2 tablespoons
1 1/4 to 1 1/2 tablespoons	2 to 3 tablespoons
6 to 7 tablespoons	8 to 12 tablespoons
3 1/4 to 1 1/4 cupfuls	1 1/4 to 2 1/4 cupfuls

THIS PACKAGE CONTAINS


Active Ingredient: Copper, expressed
as metallic, (in the form of Bordeaux
Mixture).....12.75%
Inert Ingredients.....87.25%

100.00%

CAUTION:
HARMFUL IF SWALLOWED

ACCEPTED

Aug 16, 1960
UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTER-
ED UNDER NO. 577-97



**WRITE FOR LATEST SPRAYING
AND DUSTING GUIDE**

Sold For Fungicide Purposes