

ONE POUND

NET WEIGHT

# ACME

# SEVIN

  

## 5 DUST

**Controls**  
**Mexican bean beetles,**  
**Japanese beetles,**  
**Squash bugs and certain**  
**other insect pests of**  
**fruit, tobacco, vegetables,**  
**ornamentals, and animals.**

<b>ACTIVE INGREDIENT:</b>	
Carbaryl (*1-Naphthyl N-Methyl- Carbamate).....	5.0%
<b>INERT INGREDIENTS.....</b>	<b>95.0%</b>
<b>TOTAL</b>	<b>100.0%</b>

\*Trade Mark Union Carbide Corporation for  
1-Naphthyl N-Methyl-Carbamate.  
U.S. Patent 2,903,478.

**CAUTION: KEEP OUT OF REACH OF  
CHILDREN.** See other panels for other cautions.

Made for  
**ACME QUALITY PAINTS, INC.**  
Detroit 11, Michigan Made in U.S.A.  
SOLD FOR INSECTICIDE PURPOSES

### DIRECTIONS FOR USE

Read the Cautions! Sift from this shaker top canister, or, better still, apply with a dust gun. Make application in morning or evening when it is cool. Treat upper and lower surfaces of all foliage. Do not apply in excess. Start applications at first sign of insects. Repeat as necessary, generally at 7-day intervals, oftener in rainy weather. Do not use closer to harvest than indicated following each crop. Light application should be made in seed beds if plants show signs of insects.

### FOR CONTROL OF

- Beans:** Mexican bean beetle, leafhoppers, flea beetles, corn earworm, lygus and stinkbugs.....
- Carrots:** Leafhoppers, flea beetles and spittlebugs.....
- Corn:** Corn earworm, Fall army worm, grasshoppers. (See Notes 1 & 2)...
- Cabbage and Cauliflower:** Flea beetles, cabbage worms, harlequin bugs, corn earworm.....
- Brussels Sprouts and Broccoli:** Flea beetles, cabbage worms, harlequin bugs, corn earworm.....
- Cucumbers, Melons and Squash:** Pickleworms, squash bugs, flea beetles, cucumber beetles. (See Note 3).....
- Lettuce:** Leafhoppers, army worms, stinkbugs.....
- Okra:** Stinkbugs, corn earworm.....
- Potatoes:** Colorado potato beetle, flea beetle, leafhopper.....
- Peppers:** Tomato fruitworm, flea beetles, Fall army worm, leafhoppers, lygus bug.....
- Eggplant:** Lace bugs, tomato hornworm.....
- Tomatoes:** Tomato fruitworm, flea beetle, tomato hornworm, stinkbugs.....
- Strawberries:** Meadow spittlebug, leaf roller and strawberry weevil.....

Time to Stop Use  
Before Harvest

No Limit

No Limit

Day of  
Harvest

3 Days

3 Days

Day of  
Harvest

3 Days

No Limit

No Limit

Day of  
Harvest

Day of  
Harvest

Day of  
Harvest

1 Day

### FOR CONTROL OF

Time to Stop Use  
Before Harvest

**Tobacco:** Tobacco flea beetles, apply 1 to 2 lbs. per 100 square yards when plants come up. Repeat in 10-14 days. Again just before transplanting from seed bed.....

No Limit

**Note 1.** Do not feed forage to dairy animals or animals being finished for slaughter.

**Note 2.** Apply to corn silks every 3 days until dry for earworm control.

**Note 3.** Do not use on watermelon in Florida. Do not apply to melons or cucumbers when temperatures exceed 89° F. or when vines are wet.



**Cats and Dogs:** Fleas and ticks on cats and dogs. Rub into skin, avoid eyes. Dust quarters weekly. Do not use on kittens under 2 months old.



**Chickens:** Fleas, Lice, Mites and Ticks: One pound of 5% SEVIN Dust is enough for one treatment of 100 chickens. It can also be used on pheasants, pigeons and partridge. Dust the chicken house, and roosts for mites and ticks with one pound of 5% Dust per 40 square feet. Do not dust birds within 7 days of slaughter. Avoid dusting of nests or eggs. Do not use on pet birds.

### CAUTIONS

May be harmful if swallowed or inhaled. May cause irritation of eyes, nose and throat. Avoid breathing dust. Avoid contact with eyes, skin or clothing. Wash thoroughly after use. Store away from children, pets, and foodstuffs. Remove any residues at harvest by brushing, washing or peeling. Do not apply to open blossoms; toxic to honey bees. Do not feed treated crops, pods or husks to animals being readied for slaughter.

### NOTICE

Neither the manufacturer nor the seller makes any representation or warranty, express or implied and shall not be held responsible in any manner for any personal injury, property damage, or other type of loss resulting from the handling, storage, or use of this product, whether or not used in accordance with directions.

USDA REGISTRATION NUMBER 577-462

LD-304C2 7-65 21034

11-18-65  
577-462  
295-225  
INDEX THE FOLLOWING INFORMATION FOR RECORDS TO BE KEPT UNDER THE SUBJECT TO ATTACHED COMMENTS

577-462

11/18/65

1/1