

ONE POUND

NET WEIGHT

ACME



Controls Mexican bean beetles, Japanese beetles, Squash bugs and certain other insect pests of fruit, tobacco, vegetables, ornamentals, and animals.

11-13-68 577-462

ACTIVE INGREDIENT:

Table with 2 columns: Ingredient name and percentage. Includes Carbaryl (5.0%), Inert ingredients (95.0%), and Total (100.0%).

Trade Mark Union Carbide Corporation for 1-naphthyl N-methylcarbamate. U.S. Patent 2,903,478.

CAUTION: KEEP OUT OF REACH OF CHILDREN. See left panel for other cautions.

ACME INSECTICIDE DIVISION The Sherwin-Williams Co.

8250 St. Aubin Ave., Detroit, Mich. 48211 Made in U.S.A.

SOLD FOR INSECTICIDE PURPOSES

DIRECTIONS FOR USE

Read the Cautions! Sift from this shaker top canister, or, better still, apply with a dust gun. Make application in morning or evening when it is cool. Treat upper and lower surfaces of all foliage. Do not apply in excess. Start applications at first sign of insects. Repeat as necessary, generally at 7-day intervals, oftener in rainy weather. Do not use closer to harvest than indicated following each crop. Light application should be made in seed beds if plants show signs of insects.

FOR CONTROL OF

Table listing pests and their control methods. Columns include pest name and 'Time to Stop Use Before Harvest'.

FOR CONTROL OF

Tobacco: Tobacco flea beetles, apply 1 to 2 lbs. per 100 square yards when plants come up. Repeat in 10-14 days. Again just before transplanting from seed bed. No Limit. Note 1. Do not feed forage to dairy animals or animals being finished for slaughter. Note 2. Apply to corn silks every 3 days until dry for earworm control. Note 3. Do not use on watermelon in Florida. Do not apply to melons or cucumbers when temperatures exceed 80° F. or when vines are wet.



Cats and Dogs: Fleas and ticks on cats and dogs. Rub into skin, avoid eyes. Dust quarters weekly. Do not use on kittens under 2 months old.



Chickens: Fleas, Lice, Mites and Ticks: One pound of 5% SEVIN Dust is enough for one treatment of 100 chickens. It can also be used on pheasants, pigeons and partridge. Also dust the chicken house, and roosts for mites and ticks with one pound of 5% Dust per 40 square feet. Do not dust birds within 7 days of slaughter. Avoid dusting of nests or eggs. Do not use on pet birds.

CAUTIONS

Harmful if inhaled or swallowed. Do not take internally. Avoid breathing dust or spray mist. Skin contact with any insecticide may be harmful. Avoid unnecessary contact. Repeated, prolonged, or careless use may cause cholinesterase inhibition. Wash thoroughly after use. Wear long-sleeved work clothing. Change clothing daily. Avoid contaminating feed and food-stuffs. Do not dust open blossoms. Toxic to honey bees.

Do not reuse container. Destroy when empty.

NOTE TO PHYSICIAN: Carbaryl is a moderate reversible cholinesterase inhibitor. Atropine is antidotal.

NOTICE: Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.

USDA REGISTRATION NUMBER 577-462

LD-304C7 8-68 22403

577-462

11/13/68

1/1