

ONE POUND

NET WEIGHT

ACME



Controls Mexican bean beetles, Japanese beetles, Squash bugs and certain other insect pests of fruit, tobacco, vegetables, ornamentals, and animals.

ACTIVE INGREDIENT: Carbaryl (*1-naphthyl N-methyl-carbamate) 5.0% INERT INGREDIENTS 95.0% TOTAL 100.0%

*Trade Mark Union Carbide Corporation for 1-naphthyl N-methylcarbamate. U.S. Patent 2,903,478.

CAUTION: KEEP OUT OF REACH OF CHILDREN. See left panel for other cautions.

ACME INSECTICIDE DIVISION The Sherwin-Williams Co. 8250 St. Aubin Ave., Detroit, Mich. 48211 Made in U.S.A.

SOLD FOR INSECTICIDE PURPOSES

USDA REGISTRATION NUMBER 577-462

LD-304C9 3-69 22701

DIRECTIONS FOR USE

Read the Cautions! Sift from this shaker top canister, or, better still, apply with a dust gun. Make application in morning or evening when it is cool. Treat upper and lower surfaces of all foliage. Do not apply in excess. Start applications at first sign of insects. Repeat as necessary, generally at 7-day intervals, oftener in rainy weather. Do not use closer to harvest than indicated following each crop. Light application should be made in seed beds if plants show signs of insects.

FOR CONTROL OF

- Beans: Mexican bean beetle, leafhoppers, flea beetles, corn earworm, lygus and stinkbugs. No Limit
Carrots: Leafhoppers, flea beetles and spittlebugs. No Limit
Corn: Corn earworm, Fall army worm, grasshoppers. (See Notes 1 & 2) Day of Harvest
Cabbage and Cauliflower: Flea beetles, cabbage worms, harlequin bugs, corn earworm. 3 Days
Brussels Sprouts and Broccoli: Flea beetles, cabbage worms, harlequin bugs, corn earworm. 3 Days
Cucumbers, Melons and Squash: Pickleworms, squash bugs, flea beetles, cucumber beetles. (See Note 3) Day of Harvest
Lettuce: Leafhoppers, army worms, stinkbugs. Head 3 Days Leaf 14 Days
Okra: Stinkbugs, corn earworm. No Limit
Potatoes: Colorado potato beetle, flea beetle, leafhopper. No Limit
Peppers: Tomato fruitworm, flea beetles, Fall army worm, leafhoppers, lygus bug. Day of Harvest
Eggplant: Lace bugs, tomato hornworm. Day of Harvest
Tomatoes: Tomato fruitworm, flea beetle, tomato hornworm, stinkbugs. Day of Harvest
Strawberries: Meadow spittlebug, leaf roller and strawberry weevil. 1 Day

Time to Stop Use Before Harvest

FOR CONTROL OF

Time to Stop Use Before Harvest

Tobacco: Tobacco flea beetles, apply 1 to 2 lbs. per 100 square yards when plants come up. Repeat in 10-14 days. Again just before transplanting from seed bed. No Limit
Note 1. Do not feed forage to dairy animals or animals being finished for slaughter.
Note 2. Apply to corn silks every 3 days until dry for earworm control.
Note 3. Do not use on watermelon in Florida. Do not apply to melons or cucumbers when temperatures exceed 80° F. or when vines are wet.



Cats and Dogs: Fleas and ticks on cats and dogs. Rub into skin, avoid eyes. Dust quarters weekly. Do not use on kittens under 2 months old.
Chickens: Fleas, Lice, Mites and Ticks: One pound of 5% SEVIN Dust is enough for one treatment of 100 chickens. It can also be used on pheasants, pigeons and partridge. Also dust the chicken house, and roosts for mites and ticks with one pound of 5% Dust per 40 square feet. Do not dust birds within 7 days of slaughter. Avoid dusting of nests or eggs. Do not use on pet birds.

CAUTIONS

Harmful if inhaled or swallowed. Do not take internally. Avoid breathing dust. Avoid skin contact, may be harmful. Repeated, prolonged, or careless use may cause cholinesterase inhibition. Wash thoroughly after use. Wear long-sleeved work clothing. Change clothing daily. Avoid contaminating feed and foodstuffs. Do not dust open blossoms. This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your cooperative Agricultural Extension Service.

Do not reuse container. Destroy when empty.

NOTE TO PHYSICIAN: Carbaryl is a moderate reversible cholinesterase inhibitor. Atropine is antidotal.

577-462

08/31/69

1/1