

DIRECTIONS FOR USE:

Keep lid on, shake the can prior to measuring. Apply at 1/2 lbs. per 50 gallons of water - 5 to 8 teaspoonfuls per gallon. Apply so as to completely wet all fruit and foliage. Do not apply more than 500 gallons per acre. Contains 5% actual Methoxychlor. Make application as soon as insects appear, at petal fall on fruit. Repeat at 7-14 day intervals to maintain control. Do not apply closer to harvest than shown below for specific crops.

For control of house flies, stable flies, and aid in residual control of mosquitoes, use the contents of this can in 5 quarts of water (5% actual). Apply 1 to 2 gallons per 1000 sq. ft. Run-off point. Spray around porches, windows and exterior walls of farm buildings where insects tend to congregate. Do not use this dosage on crops. Around barns, exclude all animals and cover feed and water supplies before spraying.


PRECAUTION: Wash thoroughly after handling. Avoid breathing dust while measuring. Do not feed treated crop residues to animals for feed within 7 days of harvest. Avoid contaminating feed and clothing.

Environmental Cautions: This product is toxic to fish. Keep out of lakes, streams or rivers. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water during cleaning of equipment or disposal of wastes. Apply this product in accordance with directions specified on this product label. This product may kill fish and other aquatic organisms. Avoid spraying in areas that will contaminate any body of water. Apply only as directed on this label. This product is toxic to bees and should not be applied when bees are actively visiting the area.

The Seller warrants that the product conforms to its description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use. This warranty, nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions or under abnormal conditions or conditions not reasonably foreseeable to seller and buyer assumes the responsibility of any such use.

FOR INSECTICIDE PURPOSES
DO NOT REUSE CONTAINER. DESTROY WHEN EMPTY.

Manufactured in the U.S.A. for
Herwin-Williams Co. - GREENSBORO, N.C.



***For Control of Certain Flies, Mosquitoes and Certain Chewing Insects on Shade Trees, Fruit Trees and Vegetables.**

Active Ingredient:	
Methoxychlor**, Technical	50.0%
Inert Ingredients:	50.0%
TOTAL 100.0%	

**Equivalent to 44% 2,2-bis(p-methoxyphenyl)-1,1,1-trichloroethane and 6% other isomers and reaction products.

CAUTION: Keep Out Of Reach Of Children. Read caution elsewhere on label.

ONE POUND NET WEIGHT



acme methoxychlor 50% wettable

DIRECTIONS FOR USE:

Tree Fruit:

Apples and Pears
To control codling moth, plum curculio, apple maggot, tent caterpillar and Japanese beetle. Use the stronger dosage where curculio is a problem.

Apricots, Peaches, Nectarines, Plums
To control plum curculio and Japanese beetle, use 8 tsp./gallon. Start at petal fall or when adults appear.

Cherries
To control cherry fruitworm, cherry fruit fly and rose chafer, use 5 tsp./gallon. Use 8 tsp./gallon for plum curculio and Japanese beetle.

Bush Fruit:

Blackberries, loganberries, raspberries
To control rose chafer and strawberry weevil, use 8 tsp./gallon on foliage when insects appear.

Strawberries
To control leaf tier, flea beetle, spittlebug, strawberry leaf beetle and strawberry weevil, use 8 tsp./gallon. Apply on foliage as needed.

Grapes
To control grape berry moth, grape leafhopper and rose chafer, use 5 tsp./gallon. Start at petal fall and repeat. Japanese beetle, use 8 tsp./gallon and apply as soon as insects appear.

Vegetable Gardens:

Asparagus
To control asparagus beetle, use 5 tsp./gallon. Wash to remove residues.

Beans, blackeyed peas
To control corn earworm, flea beetle, Mexican bean beetle, potato leafhopper, use 8 tsp./gallon. Do not feed vines to animals.

Beets
To control blister beetle and flea beetle, use 8 tsp./gallon. Spray foliage.

Cucumbers, squash, melons
To control cucumber beetle, use 8 tsp./gallon on foliage as soon as insects appear.

If dosage reduced to 1/2 in last two sprays.
Eggplant, peppers
To control flea beetle, use 8 tsp./gallon on foliage as needed.

Potatoes
To control leafhoppers, use 10 tsp./gallon on foliage as needed.

Tomatoes
To control blister beetle, flea beetle and tomato fruitworm, use 8 tsp./gallon. Apply on foliage as needed.
When disease is a problem, combine with Acme Zineb 75W Fungicide or Acme Wettable Dusting Sulphur.

Time to Stop Use Before Harvest

7 days

21 days

7 days

3 days

3 days

14 days

3 days

3 days

7 days (Roots)
14 days (Tops)

7 days

1 day

1 day

No Limit

7 days

DIRECTIONS FOR USE:

With the lid on, shake the can prior to measuring. Apply at 1 to 1 1/2 lbs. per 50 gallons of water — 5 to 8 teaspoonfuls per gallon. Apply so as to completely wet all fruit and foliage. Do not apply more than 500 gallons per acre — 5 lbs. actual Methoxychlor. Make application as soon as insects appear, at petal fall on fruit. Repeat at 7-14 day intervals to maintain control. Do not apply closer to harvest than shown below for specific crops.

To control house flies, stable flies, and aid in residual control of mosquitoes, use the contents of this can in 5 quarts of water (5% actual). Apply 1 to 2 gallons per 1000 sq. ft. or to run-off point. Spray around porches, windows and outside walls of farm buildings where insects tend to congregate. Do not use this dosage on crops. Around barns, remove all animals and cover feed and water supplies before spraying.

CAUTION: Wash thoroughly after handling. Avoid breathing dust while measuring. Do not feed treated crop residues to animals if sprayed within 7 days of harvest. Avoid contaminating feed and foodstuffs.

Environmental Cautions:
This product is toxic to fish. Keep out of lakes, streams or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this product.

This product may kill fish and other aquatic organisms. Avoid spraying or spillage that will contaminate any body of water. Apply only as specified on this label.

This product is toxic to bees and should not be applied when bees are actively visiting the area.


NOTICE: Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.

**SOLD FOR INSECTICIDE PURPOSES.
DO NOT REUSE CONTAINER. DESTROY WHEN EMPTY.**

Made in U.S.A. for:

The Sherwin-Williams Co. CLEVELAND, OHIO 44101

LD366 C1 10-73



33455
528