




ACCEPTED
577-1827
DEC 7 1973
UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTERED
ED UNDER NO. *Com. m. 210*



SNAIL & SLUG
KILLER

MEAL



SNAIL
&
SLUG

KILLER


MEAL

SNAIL & SLUG
KILLER

MEAL





ACCEPTED
5-17-73
DEC 7 1973
UNITED STATES DEPT. OF AGRICULTURE
FUNGICIDE AND PESTICIDE ACT
FOR ECONOMIC POISON REGISTRATION
ED UNLESS NO *Committee*



SNAIL & SLUG
KILLER

MEAL




SNAIL
&
SLUG

KILLER

SNAIL & SLUG
KILLER

MEAL






**SNAIL
&
SLUG**

KILLER

MEAL

SNAIL & SLUG KILLER

MEAL




NET WEIGHT 40 ozs. (2½ LBS.)

ACTIVE INGREDIENTS
METALDEHYDE
INERT INGREDIENTS

2.75%
97.25%
100.00%

WARNING: KEEP OUT OF REACH OF CHILDREN. SEE BACK PANEL FOR ADDITIONAL CAUTIONS.


THIS PESTICIDE MAY BE FATAL TO CHILDREN, DOGS, OR OTHER PETS IF EATEN. KEEP CHILDREN AND PETS OUT OF THE TREATED AREA. SCATTER MATERIAL. DO NOT LEAVE IN PILES. FOLLOW DIRECTIONS AND USE WITH CARE.



**SNAIL & SLUG
KILLER**

MEAL


PUSH HERE ▲ ▲ PULL OUT



ATTRACTS & KILLS SNAILS & SLUGS

K-Mart Snail & Slug Killer is a special preparation for the control of snails. K-Mart Snail & Slug killer attracts and kills snails and slugs with Metaidehyde. The presence of snails and slugs can be detected by their shiny mucuous like trail. They are night feeders and prefer damp, cool places in the ornamental garden, flowers, under boards or flower pots and around fences, hedges and bushes.

NOTE: This package was full when packed. Contents are sold by weight.



SNAIL & SLUG KILLER



DIRECTIONS

Water area to be treated thoroughly. Broadcast K-Mart Snail & Slug Killer late in the day in the areas likely to be infested. Apply to the soil and not directly on plants. Scatter, do not leave in piles.

Continuous watering and high temperatures cause this bait to gradually lose some of its effectiveness. Repeat applications every week or ten days during spring and summer months. Treatment every two weeks during fall and winter months is usually satisfactory. Repeated treatments provide control of new broods emerging during this period. Three baitings ten days apart, usually gives control of new hatch.

Apply one pound of meal to each 250-300 square feet.

WARNING

MAY BE FATAL IF SWALLOWED. Harmful dust. Avoid inhalation of dust. Avoid contact with skin, eyes, and clothing. Avoid food contamination. Wash hands with soap and water after using.

IF SWALLOWED, INDUCE VOMITING. CALL A PHYSICIAN IMMEDIATELY. Keep away from children, pets, and livestock. Keep animals and people off treated areas. This bait is attractive to deer and turkeys. Keep off plants. Do not reuse container. Destroy when empty.

This product is toxic to wildlife. Birds feeding on treated areas may be killed.

MANUFACTURED IN U.S.A. FOR **SK** S. S. KRESGE COMPANY, DETROIT, MICHIGAN 48232
BY SWIFT CHEMICAL COMPANY, CHICAGO, ILL. 60604

E.P.A. Reg. No. 5571624 ZA 12000