CAUTION

DDX contains as its active ingredients an anticoagulant which, if taken by humans, domestic and wildlife animals and pets may reduce the clotting ability of the blood and a serious homorrhage may result. Baits should be placed in areas inaccessible to children, pets, wildlife and domestic animals or in tamper proof bait boxes.

ANTIDOTE

In case baits are eaten give tablespoonful of salt in glass of warm water and repeat until vomit fluid is clear. Call physician immediately.

NOTE TO PHYSICIANS

When a human has known to have ingested DDX, blood transfusions combined with intravenous injections and oral doses of Vitamin K are indicated as in the case of hemorrhage caused by overdoses of dicumarol.

GUARANTEE OF SATISFACTION

Satisfaction Guaranteed or Money Refunded. If DDX does not live up to this guarantee, return the empty package to B. E. Miller Co., Hancock, Md. for full refund.

-



DIRECTIONS

Place 4 oz. of DDX in locations froquanted by rats and mice and protect from children, dogs, cats, and livestock by means of bait boxes. Baits should be replaced as consumed. Permanent bait stations should be set up if you are near a location that is infested with rats and watch from time to time to see If there is sufficient bait left. Spoiled bait should be replaced promptly.

RAT CONTROL

Four ounces of bait per bait location should be minimum amount placed. An uninterrupted supply of bait should be maintained for approximately 10 days.

MOUSE CONTROL

Place tablespoon amounts of bait at 8-12 foot intervals. An uninterrupted supply of bait should be maintained for at least 15 days.

USDA Registration Nc. 584-5 EPA Est. No. 00584

NET WT. 1 LB.

ACCEPTED JUL 20 1974

FUNG CALL TIME ROLL. 2011 A 1 1 1 1