[212.04]

				L	L,,
	5-	7-	• 7	V	1
		5-0			
, (~ 7			-	

ACTIVE INGREDIENTS:

Rotenone.	1.00%
Other Cube Resins.	2.15%
INERT INGREDIENTS	96.85%
TOTAL 1	00.00%

ASPARAGUS — Asparagus Beetles

BEAN — Mexican Bean Beetle, Bean Leaf Beetle and Pale Striped Cucumper Beetle.

CABBAGE — Imported Cabbageworm, Cabbage Looper and Spotted Cucumber Beetle.

PEPPER, SPINACH AND TOMATO — Potato Flea Beetle.

DIRECTIONS

Apply at first sign of infestation at a rate of 15 to 40 pounds per acre, depending on the size and spacing of the plants and the method of application. Repeat every 7 to 10 days or after as needed. Do not use on crops within one day of harvest.

CAUTION: KEEP OUT OF REACH OF CHILDREN

Avoid inhalation of dust.

Do not re-use empty container. Destroy it by burying with waste or burning. Stay away from smoke or fumes.

This product is toxic to fish. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

NON-WARRANTY

Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

E.P.A. REG. NO. 595-69