

DICHLONE DUST NO. 2

FUNGICIDE - CONTAINS 2% PHYGON®

ACTIVE INGREDIENTS:

Dichlone (2, 3-Dichloro-1, 4-Naphthoquinone)	2.0%
INERT INGREDIENTS:	98.0%
Total	100.0%

Dichlone Dusts are recommended for the control of certain fungus diseases on Apples, Cherries, Peaches and Plums. It can be applied by airplane or ground equipment. If it is desirable Dichlone Dusts may be mixed with sulfur. Contact your Local State Experiment Station or Extension Service for other approved formulations available in your area.

CAUTION:

MATERIAL CAUSES EYE OR SKIN IRRITATION

Keep Out Of Reach Of Children

Avoid inhalation or contact with skin or eyes. In case of contact with skin wash thoroughly with soap and water; for eyes, flush with plenty of water for at least fifteen minutes and get medical attention. Wash hands and face with soap and water after using and before eating or smoking. Avoid contamination of feed and foodstuffs.

Toxic to fish. Do not contaminate any body of water nor apply when conditions favor drift from the crop area treated. Do not contaminate water by cleaning of equipment, or disposal of wastes.

Do not reuse empty container. Destroy it by burying with waste or burning. Stay away from smoke or fumes.

NON-WARRANTY

Seller makes no warranty, expressed or implied, concerning the use of this product, other than indicated on the label. Buyer assumes all risk of use and/or handling, when such use and/or handling is contrary to label instructions.

DIRECTIONS FOR USE

APPLES – For Control of Apple Scab from Delayed Dormant through the Early Cover Sprays, apply 2% Dichlone Dust on a protectant schedule at the rate of 45 to 60 lbs. per acre before, during, or after a rain starts up to 50 hours. When 2% Dichlone Dust is applied on a protectant schedule, because of its combined protectant and eradicator properties, it is not necessary to dust more often than once in every 5 days.

CHERRIES, PEACHES and PLUMS – For Control of Brown Rot Blossom Blight of Stone Fruits, apply 2% Dichlone Dust at the rate of 45 to 60 lbs. per acre at delayed dormant, pre-pink, pink, full bloom and petal fall or shuck fall. When cool, wet weather occurs, resulting in a prolonged bloom period, apply dust every 4 days.

For the control of Brown Rot of Fruit on Cherries and Plums (Fresh Prunes), apply 2% Dichlone Dust at the rate of 45 to 60 lbs. per acre at 7 to 10-day intervals beginning at petal fall or shuck fall and continuing until just before harvest. **DO NOT APPLY WITHIN 3 DAYS OF HARVEST.**

For control of Pre-harvest Brown Rot of Peach, apply 2% Dichlone Dust at the rate of 45 to 60 lbs. per acre at 14 and 7 days before harvest.

Seller makes no warranty, expressed or implied, concerning the use of this product, other than indicated on the label. Buyer assumes all risk of use and/or handling, when such use/or handling is contrary to label instructions.

PHYGON® is the Reg. Trademark of Naugatuck Chemical Division of United States Rubber Company.

USDA Reg. No. 595-177

Net Wt. 50 Lbs.

ACCEPTED
AUG 4, 1971
UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTRATION
ED UNDER NO. 595-177



1845 STERLING N W

GRAND RAPIDS MICHIGAN