

## (FUNGICIDE DUST)

### ACTIVE INGREDIENTS:

Maneb (Manganous ethylene bisdithiocarbamate) ..... 7.0%

INERT INGREDIENTS: ..... 93.0%

100.0%

### DIRECTIONS

Haviland 7 M Dust should be applied at the rate of 15 to 30 lbs. per acre depending on the size and spacing of the plants and the method of application. In general, the lower rates are usually used early in the season and gradually increased as the foliage and plants increase in size with the higher rates being used later in the season as plants reach maturity.

**BEANS** – To control Downy Mildew and Rust – apply dust when disease first appears and repeat at weekly intervals. Do not apply to stringbeans within 4 days of harvest, no time limitation on dry beans.

**CABBAGE, BRUSSEL SPROUTS, CAULIFLOWER, KOHLRABI and BROCCOLI** – To control Downy Mildew and Alternaria Leaf Spot begin dusting as soon as disease threatens and repeat at 7 to 10 day intervals in the field. Use at 3 day intervals in plant beds. Remove residues by trimming and washing. Do not apply within 7 days of harvest on cabbage.

**CARROTS** – To control Leaf Blight and Cercospora Leaf Spot apply dust when diseases are reported in the area and continue at 7 to 10 day intervals during the growing season, or as required for adequate control.

**CELERY** – To control Early and Late Blight begin dusting in the plant bed when plants first come through the ground. Make further applications at 3 to 5 day intervals until plants are set in the field. Continue field applications on regular weekly schedule. Remove residues by stripping, trimming and washing.

**CANTALOUPE, CUCUMBER, SQUASH, MELONS and WATERMELONS** – To control Downy Mildew, Anthracose and Gummy Stem Blight start dusting when runners begin to form or at the first sign of disease. Repeat at weekly intervals. Dusting of the underside of leaves is necessary for good control.

**EGGPLANTS and PEPPERS** – To control Fruit Rot and Anthracnose apply dust as fruit begins to form and repeat weekly until harvest.

**ENDIVE, LETTUCE and SPINACH** – To control Downy Mildew and White Rust apply dust at the first signs of disease and repeat at 7 to 10 day

intervals until harvest as required. Remove residues by trimming and washing. Do not use within 7 days of harvest.

**TOMATOES** – To control Early and Late Blight, Anthracose, Stemphylium, Septoria Leaf Spot and Cladosporium Gray Mold on tomatoes in the field, greenhouse or plant beds begin dusting when disease is first reported in the area and repeat at 7 to 10 day intervals or oftener during severe late blight conditions. Use at the rate of 40 lbs. per acre in severe anthracnose areas. To avoid damage, do not use on tender young plants under glass.

**ONIONS** – To control Downy Mildew, Botrytis Leaf Blight and Purple Blotch apply dust when diseases are reported in the area and continue at 7 to 10 day intervals during the growing season.

**POTATOES** – To control Early and Late Blight apply dust at 7 to 10 days throughout the season. Begin applications early using the lower rate and increase amount per acre as vines increase in size.

### CAUTION: Keep Out of Reach of Children

Maneb Dust may cause irritation of nose throat, eyes and skin. Do not breathe dust or spray mist. In case of skin contact, wash with water; for eyes, get medical attention. **Important:** Keep dust away from fire and sparks. Store in cool dry place. Do not allow to become wet or overheated in storage. This may bring on chemical changes which will impair the fungicidal effectiveness of the dust and may generate flammable vapors. Keep container closed when not in use.

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment, or disposal of wastes.

Do not reuse empty container. Destroy it by burying with waste or burning. Stay away from smoke or fumes.

### NON-WARRANTY

Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

USDA Reg. No. 595-180

Net Wt. 50 Lbs.



# HAVILAND

1845 STERLING, N.W. GRAND RAPIDS, MICHIGAN 49502