PERTHANE MALATHION DUST NO. 64

ACTIVE INGREDIENTS:

| Diethyl Diphenyl Dichloroethane and | 5.7 % |
|---|---------------|
| Related Compounds | $0.3^{\rm o}$ |
| Malathion (O,O-Dimethyl dithiophosphate | |
| of diethylmercaptosuccinate) | 4.0% |
| INERT INGREDIENTS: | 90.0% |
| TOTAL | 100.0% |

CAUTION: Keep Out of Reach of Children

Avoid breathing dust. Avoid contact with skin. Wash thorough ly with soap and water after handling and before eating or smoking. Avoid contamination of feed and foodstuffs.

Do not reuse empty container. Destroy it by burying with wastes or burning. Stay away from smoke or fumes.

This product will kill fish and wildlife. Do not contaminate any body of water nor apply when conditions favor drift from the area treated. Birds feeding on treated areas may be killed. Do not contaminate water by cleaning of equipment or disposal of wastes.

NOTE TO PHYSICIAN: Malathion upon repeated or prolonged contact may cause Cholinesterase inhibition. Atropine is the antidote.

NON-WARRANTY

Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

DIRECTIONS

The following directions are furnished by us and are believed to be reliable. State Agricultural Experiment Station should be consulted in each agronomic area for specific timing and rates of application according to severity of local insect conditions. For general usage on crops listed below, apply 25 to 40 lbs. of this material per acre at first signs of insects or insect damage and repeat at weekly intervals as needed.

BROCCOLI and BRUSSELS SPROUTS - To control Aphids, Cabbage Looper and Imported Cabbage Worms. Do not apply within 3 days of harvest on broccoli or within 7 days of harvest on brussells sprouts.

CABBAGE AND CAULIFLOWER - To control Aphids, Cabbage Looper, Diamondback moths and Imported Cabbage Worms. Do not apply within 7 days of harvest.

KOHLRABI To control Aphids, Cabbage Loopers and Imported Cabbage Worms. Do not apply within 7 days of harvest.

LETTUCE - To control Aphids, Cabbage Loopers, Imported Cabbage Worms, Leafhoppers, and Mites. Do not apply to leaf lettuce within 14 days of harvest or to head lettuce within 7 days of harvest.

SPINACH To control Aphids, Cabbage Loopers, and Leaf hoppers. Do not apply within 7 days of harvest.

USDA Reg. No. 595-203 8-30-71 Net Wt. 50 Lbs.



HAVILAND

1845 STERLING, N.W. GRAND RAPIDS, MICHIGAN 49502