

COPPER SULFUR DUST No. 720

Contains 7% Metallic Copper and 20% Sulfur

ACTIVE INGREDIENTS	
Metallic Copper (from Triplex Copper Sulfate)	7.0
Sulfur	20.0
INERT INGREDIENTS	73.0
Total	100.0

Recommended for control of many fungal diseases on Vegetable and Tree Crops and on Dairy, Mulberry and Grape.

DIRECTIONS

For directions on application, see label. Apply at the rate suggested below depending on the crop and disease to be controlled and the method of application. Repeat weekly, if necessary, until the disease is controlled.

CARROTS To control Early Blight, apply 35 to 50 lbs. per acre weekly. For control of Late Blight, apply 35 to 50 lbs. per acre weekly. For control of Purple Blight, apply 35 to 50 lbs. per acre weekly.

CELERY To control Early Blight, apply 35 to 50 lbs. per acre weekly. For control of Late Blight, apply 35 to 50 lbs. per acre weekly. For control of Purple Blight, apply 35 to 50 lbs. per acre weekly.

BEETS To control Early Blight, apply 35 to 50 lbs. per acre weekly. For control of Late Blight, apply 35 to 50 lbs. per acre weekly. For control of Purple Blight, apply 35 to 50 lbs. per acre weekly.

CUCUMBERS, SQUASHES and MELONS To control Early Blight, apply 35 to 50 lbs. per acre weekly. For control of Late Blight, apply 35 to 50 lbs. per acre weekly. For control of Purple Blight, apply 35 to 50 lbs. per acre weekly.

TOMATOES, EGG PLANT and PEPPERS To control Early Blight, Late Blight, Septoria Blight, Phomopsis Blight and Anthracnose, apply 35 to 50 lbs. per acre. Begin first application in seed beds and continue weekly until harvest.

POTATOES To control Late Blight, apply 35 to 50 lbs. per acre weekly. Begin first application after emergence and repeat at weekly intervals until harvest.

GRAPES and SPINACH To control Downy Mildew, apply 35 to 50 lbs. per acre. Begin application seven to ten days after foliage appears and repeat weekly as long as necessary to control disease.

ONIONS To control Downy Mildew and Purple Blight, apply 35 to 50 lbs. per acre. For Purple Blight, use Zineb. Make first application 10 to 30 days after plants come up and repeat at intervals.

CAUTION: Keep Out of Reach of Children

Keep out of reach of children. Do not feed to animals. **KEEP OFF SWALLOWED**

See formula, burn foliage. Do not use on plants with frost damage.

8-16-71

NON-WARRANTY

USE OF THIS PRODUCT IS AT THE USER'S SOLE RISK. THE USER ASSUMES ALL LIABILITY FOR ANY DAMAGE TO PERSONS OR PROPERTY. THE USER AGREES TO HOLD THE MANUFACTURER HARMLESS FROM ALL SUCH LIABILITY.

USDA Reg. No. 595-204

Net Wt. 50 Lbs



1845 STERLING, N.W. GRAND RAPIDS, MICHIGAN 49502