

ACTIVE INGREDIENTS:

dodine (dodecylguanidine acetate) 2.0%

INERT INGREDIENTS: 98.0%

Total 100.0%

A FUNGICIDE DUST FORMULATION SUGGESTED FOR USE BY GROUND DUSTER OR AIRPLANE, FOR CONTROL OF APPLE AND PEAR SCAB.

CAUTION:

CONCENTRATED MATERIAL CAUSES EYE IRRITATION

Keep Out Of Reach Of Children

In case of contact with eyes, flush eyes with plenty of water for at least fifteen minutes and get medical attention.

RECOMMENDATIONS FOR USE

APPLES: Apple Scab – Apply 40 lbs. of 2% Dodine Dust per acre beginning with first cover application as needed. Use 25 to 40 lbs. of a 3% or 4% Dodine Dust per acre at 5-7 day intervals or as needed to maintain scab control from pre-bloom to first cover.

Do not apply within 5 days of harvest.

TREATED apples should not be used in the manufacture of apple pomace for use in livestock feeds.

USDA Reg. No. 595-219

PEARS: Pear Scab Apply 40 lbs. of 2% Dodine Dust per acre beginning with first cover application as needed. Use 25 to 40 lbs. of 3% or 4% Dodine Dust per acre at 5-7 day intervals or as needed to maintain scab control from pre-bloom to first cover.
Do not apply within 5 days of harvest.

ADDITIONAL CAUTIONS

Do not apply dodine in combination with Lime or Chlorobenzilate.

Avoid drift of dust on adjacent crops.

Care should be taken to avoid application on peaches, plums and bearing sweet cherries since severe foliage or fruit injury may result.


Do not graze cover crops in treated orchards.

Consult your State Agricultural Experiment Station or Extension Service for proper timing of applications.

NON WARRANTY

Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instruction.

CYPREX® is the registered trademark of the American Cyanamid Co.



Net Wt. 50 Lbs.

ACCEPTED WITH COMMENTS



HAYILAND

1845 STERLING, N. W.

GRAND RAPIDS, MICHIGAN