

(CONTAINS 7% METALLIC COPPER)

**ACTIVE INGREDIENTS:**

Metallic Copper (From Tribasic Copper Sulfate) .....	7.0%
<b>INERT INGREDIENTS:</b> .....	<b>93.0%</b>
Total	100.0%

Recommended for control of many fungus diseases on Vegetable and Truck Crops and for Downy Mildew on Grapes.

**DIRECTIONS**

The following directions are furnished by us and are believed to be reliable but in no way guaranteed. This dust should be applied at the rates suggested below depending on the size and spacing of the plants and the method of application.

**CARROTS** – To control Alternaria Blight, apply 45 lbs. per acre after emergence and repeat at 7 to 10 day intervals or after a rain if needed.

**CELERY** – To control Early and Late Blight, apply 30 to 50 lbs. per acre after emergence and repeat at 7 to 10 day intervals. For best control - alternate application with Haviland Maneb or Thiram Dust.

**BEETS** – To control Leaf Spot, apply 35 to 50 lbs. per acre at 7 to 10 day intervals after emergence.

**CUCUMBERS, SQUASHES and MELONS** – To control Angular Leaf Spot, Alternaria Leaf Blight, Scab, and Anthracnose, apply 30 to 50 lbs. per acre weekly after emergence. For Alternaria Leaf Blight, a mixture of 25 lbs. of Copper Fungicide Dust No. 7 with 30 lbs. of a 7% Zineb dust will give better results.

**TOMATOES, EGGPLANT and PEPPERS** – To control Early Blight, Late Blight, Septoria Blight, Phomopsis Blight and Anthracnose, apply 35 to 50 lbs. per acre. Application should begin in the seedbed and continued at 7 day intervals in the field.

**POTATOES** – To control Late Blight, apply 35 to 50 lbs. per acre weekly after emergence.

**GRAPES AND SPINACH** -- To control Downy Mildew, apply 35 to 50 lbs. per acre early in the growing season or after emergence and repeat at 7 to 10 day intervals.

**ONIONS** – To control Downy Mildew and Purple Blotch, apply 35 to 50 lbs. per acre with 6 to 10 lbs. of sulfur or use a 7% Copper, 20% Sulfur Dust. This should be alternated with 10% Maneb or Thiram. For Purple Blotch use Zineb in place of Thiram. Applications should begin as soon as plants come up and repeated at 7 to 10 day intervals.

**NON-WARRANTY**

Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling when such use and/or handling is contrary to label instructions.

**CAUTION: Keep Out of Reach of Children**

Harmful if Swallowed!  
Avoid inhalation of dust.  
Do not contaminate Feed or Foodstuffs.

8-4-71  
595-221

USDA Reg. No. 595-221

Net Wt. 50 Lbs.

Comments



1845 STERLING N.W. GRAND RAPIDS, MICHIGAN 49502

ACCEPTED WITH COMMENTS