

HAVILAND CAPTAN-STREPTOMYCIN 7.5-.01

GUARANTEE

ACTIVE INGREDIENTS:

Captan (N-Trichloromethylthio-4-cyclohexene-1, 2-dicarboximide)	7.50%
Streptomycin01%
INERT INGREDIENTS:	92.49%

100.00%

Store in Cool Dry Place – Protect from Excessive Heat

RECOMMENDATIONS

This material is especially formulated for potato seed piece treatment.

POTATO SEED PIECES: To control Seed Rot and Damp-Off, Verticillium Wilt, and Brown Eye Disease, dust at 10 ounces per bushel of cut seed pieces (equivalent to 1 lb. per 100 lbs. of cut seed pieces). This material may be applied in any

Net Wt. 50 lbs.

convenient container or by duster attachment mounted over conveyor belt. Dust treatment should be made within 6 hours of cutting. When planting is to be delayed for more than 1 or 2 days, the treated pieces should be stored in open crates for 2 or 3 days before bagging.

NON-WARRANTY

Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instruction.

CAUTION: Keep Out of Reach of Children

- May cause skin irritation.
- Avoid contact with skin and eyes.
- Avoid breathing dust.
- Wash with soap and water after exposure.
- Avoid contamination of feed and foodstuffs.

USDA Reg. No. 595-227



HAYILAND

1845 STERLING N W

GRAND RAPIDS MICHIGAN