

MALATHION COPPER DUST

[595-229]

ACTIVE INGREDIENTS

Malathion (O,O-dimethyl dithiophosphate of diethyl mercapto succinate)	5.0%
Copper (metallic form) (basic Copper Sulfate)	7.0%
INERT INGREDIENTS	88.0%
Total	100.0%

ACCEPTED
 SEP 2 1971
 UNDER THE FEDERAL INSECTICIDE
 FUNGICIDE AND RODENTICIDE ACT
 FOR ECONOMIC POISON REGISTERED
 UNDER NO. 595-229
 TO ATTACHED COMMENTS.

DIRECTIONS

The following directions are furnished by us and are believed to be reliable. Make thorough, full-coverage applications as insects or disease threatens. Repeat applications at 5 to 7 days as necessary. Rates suggested below are in lbs. per acre. Do not use excessive amounts or closer to harvest than recommended below. Consult your State Agricultural Authorities for proper timing of applications.

Crop	Insect	Disease	Rate	Limitations
BEANS	Mexican Bean Beetle	Angular leaf spot	30-35	1 day
	Leafhoppers	Anthrachnose		
	Spider Mites	Downy mildew		
BEETS	Aphids	Leaf blight	40-45	7 days
		Leaf spot		
CABBAGE, CAULIFLOWER, and BROCCOLI	Imported Cabbageworm	Downy mildew	40-45	7 days*
	Cabbage looper	Leaf spot		
	Aphids			
	*May be used on broccoli to within 3 days.			
CARROTS	Aphids	Early & late blight	25-30	7 days
	Leafhoppers	Leaf Spot		
CUCUMBERS, MELONS, and SQUASH	Aphids	Angular leaf spot	30-40	1 day
	Cucumber beetles	Anthrachnose		
	Pickleworm			
	Spider Mites			

TOMATOES, EGGPLANT, and PEPPERS

Aphids	Early & late blight	25-30	3 days
Spider mites	Septoria leaf spot		
Drosophila	Leaf Mold		

ONIONS POTATOES

Thrips	Downy mildew	25-30	3 days
Aphids	Early & late blight	30-45	0
Leafhoppers			

CAUTION: Keep Out of Reach of Children

Harmful if swallowed. Avoid prolonged breathing of dust. Avoid repeated or prolonged contact with skin. For eyes, wash with plenty of water for 15 minutes and get medical attention. Wash thoroughly after handling and before eating or smoking. Avoid contamination of feed and foodstuffs. Do not allow product to become wet in storage. This may lead to a reduction of effectiveness of product.

Highly toxic to fish. Do not contaminate any body of water nor apply when conditions favor drift from the crop area treated.

Do not reuse empty container. Destroy it by burying with waste or burning. Stay away from smoke or fumes.

NOTE TO PHYSICIANS: Upon repeated, long use, Malathion may cause Cholinesterase inhibition. Atropine is the antidote.

NON-WARRANTY

Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling when such use and/or handling is contrary to label instructions.

USDA Reg. No. 595-229

Net Wt. 50 Lbs.



HAYILAND

1845 STERLING, N.W. GRAND RAPIDS, MICHIGAN 49502

ACCEPTED WITH COMMENTS