

[595-234]

597
 5-3.5-3.5
 (27)

MALATHION MANEB COPPER DUST NO. 5-3.5-3.5

ACTIVE INGREDIENTS:

Malathion (O,O dimethyl dithiophosphate of diethyl mercaptacetate)	5.0
Maneb (Manganoous ethylene bisdithiocarbamate)	3.5
Elemental Copper (from Tribasic Copper Sulfate)	3.5
INERT INGREDIENTS	88.0%
Total	100.0%

Recommended for control of various insects and fungal diseases on most vegetables and truck crops. Specific recommendations may vary in different agricultural areas. It is suggested that you consult your State Agricultural Authorities before using this material. If crop to be dusted is for canning purposes, consult your primary fieldman before applying dust.

CAUTION: Keep Out of Reach of Children

Harmful if swallowed. Avoid prolonged breathing of dust. Avoid repeated or prolonged contact with skin. If eye is irritated with plenty of water for 15 minutes and get medical attention if irritation thoroughly after bandaging and before eating or smoking. Avoid contamination of food and face stuff. Do not allow product to become part of animal's forage. This may lead to a reduction in effectiveness of product. **NOTE TO PHYSICIANS:** Upon repeated use, Malathion may cause Cholinesterase inhibition. Antidote is the atropine.

Do not get dust on or contaminate any body of water. Do not apply when conditions are such that the dust will be carried.

Always use empty containers. Destroy if by burning with a ste. or cutting far away from stake or house.

DIRECTIONS

The following directions are furnished by us and are believed to be of value. Apply dust at the rate of 25 to 30 lbs. per acre. Use when insects appear or disease threatens. Repeat application weekly as necessary, depending on the severity of the infestation.

CROP	INSECTS	DISEASE	DAYS BETWEEN LAST APPLICATION AND HARVEST
BEANS	Mexican Bean Beetle Leafhopper Spider Mites	Rust Anthracnose Downy Mildew	1 (dry) 3 (succulent)
CARROTS & CELERY	Aphids Spider Mites	Cercospora Leaf Spot Early & Late Blight	7 (Remove excess residues on celery by stripping, trimming and washing)
* CUCUMBERS, MELONS & SQUASH	Aphids Spider Mites Leafhopper	Anthracnose Angular Leaf Spot Downy Mildew	1
ONIONS	Thrips	Purple Blotch Downy Mildew	3
POTATOES	Aphids Leafhopper	Early & Late Blight	0
TOMATOES, EGGPLANT & PEPPERS	Aphids Spider Mites	Early & Late Blight Anthracnose Septoria Gray Leaf Spot	1 (Tomatoes) 3 (Eggplant & Peppers)

*Do not apply this dust to cucurbits unless plants are dry.

NON-WARRANTY

Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling, when such use and/or handling is contrary to label instructions.

USDA Reg. No. 595-234

Net Wt. 50 Lbs.



1845 STERLING, N.W. GRAND RAPIDS, MICHIGAN 49502