

# Dichlone Sulfur Dust No. 2-20

## ACTIVE INGREDIENTS:

Dichlone (2, 3-Dichloro-1, 4-napthoquinone) .....	2.0%
Sulfur .....	20.0%
INERT INGREDIENTS: .....	78.0%
	100.0%

Recommended for the control of fungus disease on apples, cherries, plums and peaches. This dust is compatible with the dry and wettable powders of DDT, Rotenone, Chlordane, Aldrin, Dieldrin, Parathion, Aramite, Thiram, Ferbam, Captan and Zineb.

## DIRECTIONS

The following directions are furnished by us and are believed to be reliable and of value. For further information consult your state experiment station for recommended dosages in your area.

**APPLES:** To control Apple Scab and Powdery Mildew, apply this dust at the rate of 45 to 60 lbs. per acre. Begin applications at the first sign of green tissues and repeat applications every 5 to 7 days up through the first cover spray (7 days after petal fall). Some browning of petals may occur, but this will not discourage bees or affect fruit set. Do not use this dust after first cover.

**PEACHES, CHERRIES AND PLUMS:** To control Peach Scab, Brown Rot and Leaf Spot, apply this dust at the rate of 45 to 60 lbs. per acre. Applications should begin at the pre-bloom stage and continue every 5 to 7 days through the bloom, petal fall and first four cover applications. Dust may be used every 3 to 4 days if rainy weather prevails. For Brown Rot and Leaf Spot, this dust may be used at weekly intervals to within 7 days of harvest on Peaches and to within 3 days of harvest on Cherries and Plums.

UNDER THE FEDERAL INSECTICIDE  
FUNGICIDE AND RODENTICIDE ACT,  
FOR ECONOMIC POISON REGISTRATION  
ED UNDER NO.

NO WARRANTY

*Comment*

Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

**CAUTION:** Keep Out of Reach of Children

May cause skin irritation. Avoid prolonged contact with skin. Avoid inhalation of dust. Wear protective clothing. Wash with soap and water after using and before eating or smoking. Avoid storage near feed or foodstuffs.

Birds feeding on treated areas may be harmed.

Sulfur may burn foliage when temperatures are exceedingly high.

Do not apply when weather conditions favor drift of dust from crop area treated.

USDA REG. NO. 595-237

NET WT. 50 LBS.



**HAYILAND**  
AGRICULTURAL CHEMICAL COMPANY

1845 STERLING N. W.

GRAND RAPIDS, MICHIGAN

ACCEPTED WITH COMMENTS