

[395-240]

MALATHION MANEB DUST NO.57

ACTIVE INGREDIENTS:

Malathion (O,O dimethyl dithiophosphate of diethyl mercaptosuccinate)	5.0%
Maneb (Manganous ethylene bisdithiocarbamate)	7.0%
INERT INGREDIENTS:	88.0%
TOTAL	100.0%

CAUTION: Keep Out of Reach of Children

Harmful if swallowed. Avoid prolonged breathing of dust. Avoid repeated or prolonged contact with skin. For eyes, wash with plenty of water for 15 minutes and get medical attention. Wash thoroughly after handling and before eating or smoking. Avoid contamination of feed and foodstuff. Do not allow product to become wet during storage. This may lead to a reduction of effectiveness of product.

This product is toxic to fish. Keep out of lakes, streams, or ponds. Do not apply when weather conditions favor drift from area treated.

Do not reuse empty container. Destroy it by burying with waste or burning. Stay away from smoke or fumes.

Note to physician:

Upon repeated, long use, Malathion may cause Cholinesterase inhibition. Atropine is the antidote.

Recommended for control of various insects and fungus diseases on certain vegetables and truck crops. As specific recommendations may vary in different agricultural areas, it is suggested that you consult your State Agricultural Experiment Station before using this material. If crop to be dusted is for canning purposes, consult your cannery fieldman before applying dust.

USDA Reg. No. 595-240

Net Weight 50 Lbs.

DIRECTIONS

The following directions are furnished by us and are believed to be of value. Apply dust at the rate of 25 to 30 lbs. per acre when insects appear or disease threatens. Repeat application weekly as often as necessary, depending on the severity of the infestation.

Crop	Insects	Disease	Days Between Last Application and Harvest
BEANS	Mexican Bean Beetle	Rust	1 (dry)
	Leafhopper	Anthracnose	4 (succulent)
	Spider Mites	Downy Mildew	
*CARROTS & CELERY	Aphids	Cercospora leaf spot	7
	Spider Mites	Early & late Blight	
**CUCUMBERS, MELONS and SQUASH	Aphids	Anthracnose	1
	Spider Mites	Downy mildew	
	Leafhopper		
ONIONS	Thrips	Purple Blotch	3
		Downy Mildew	
POTATOES	Aphids	Early & Late Blight	0
	Leafhopper		
TOMATOES, EGGPLANT and PEPPERS	Aphids	Early & Late Blight	3
	Spider Mites	Anthracnose	
		Septoria	
		Gray Leaf Spot	

*Remove excess residue by stripping, trimming and wash
 **Do not apply this dust to cucurbits unless plants are dry.

NON-WARRANTY

Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.



1845 STERLING, N.W. GRAND RAPIDS, MICHIGAN 49502

REGISTERED WITH COMMENTS