

**RESTRICTED USE PESTICIDE**

**PARATHION COPPER NO. 1.5-7**

<b>ACTIVE INGREDIENTS</b>	
Parathion (O,O Diethyl O-p-trophenyl phosphorothioate)	1.5%
Copper metal (from Tribasic Copper Sulfate)	7.0%
<b>INERT INGREDIENTS</b>	91.5%
<b>Total</b>	<b>100.0%</b>

Highly toxic to fish and wildlife. Do not contaminate streams, lakes, or ponds, nor apply when conditions favor drift from treated area. Do not apply to any area not specified on the label.

**DIRECTIONS**

As recommendations for the use of Parathion products vary in different agricultural areas it is suggested that you consult your State Agricultural Experiment Station before using this product. If the crop to be dusted is for canning purposes, consult your cannery fieldman before applying dust. The following general directions are furnished by us and are believed to be of value but in no way guaranteed.

**CARROTS** To control Aphids, Six Spotted Leafhopper and Carrot Blight apply dust at the rate of 20 to 35 lbs. per acre depending on the size and spacing of the plants. Begin application early when plants are coming up and repeat applications at weekly intervals as long as insects or blight threatens. Do not apply within 15 days of harvest. Do not use treated carrot tops for food or feed. For adequate control of Leafhoppers it is necessary to treat margins of fields also.

**CELERY** To control Aphids, Six Spotted Leafhoppers, Tarnish Plant Bugs, Early and Late Blight apply dust at the rate of 20 to 30 lbs. per acre at weekly intervals. Begin applications as soon as insects appear or blight threatens. Do not apply within 21 days of harvest.

**CUCUMBERS, MELONS and SQUASH** To control Cucumber Beetle, Aphids, Plant Bugs, Melonworm, Pumpkinworm, Angular Leaf Spot, Alternaria Disease, Scab and Anthracnose apply dust at the rate of 20 lbs. per acre when insects appear or disease threatens. Do not apply within 15 days of harvest in cucumbers. Do not use within 15 days of harvest in melons and within 7 days of harvest in squash.

**POTATOES** To control grasshopper and late blight apply when insect increases in number and late blight affected as regular spray or dust schedule or when blight threatens. Use at the rate of 20 to 30 lbs. per acre. Do not apply within 5 days of harvest.

**NON-WARRANTY**

We make no warranty, expressed or implied, concerning the use of this product. The user assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

E.P.A. Reg. No. 595-241  
E.P.A. Est. No. 10183-MI-4

Net Wt. 50 Lbs.

INTERNET

EXPIRES

**DANGER: Keep Out Of Reach Of Children**


**POISONOUS IF SWALLOWED OR INHALED OR ABSORBED THROUGH SKIN, OR EYES RAPIDLY ABSORBED THROUGH SKIN.**

Do not get in eyes, clothing or on skin. In case of contact, wash immediately with soap and water. Wear natural rubber gloves, protective clothing and goggles. Wear mask or respirator of a type approved by the U.S. Bureau of Mines for parathion protection. Wash arms, hands and face thoroughly after handling and before eating or smoking. Wear clean clothes daily and wash contaminated clothing with soap and hot water before reuse. Keep away from children, animals, feed and feedstuff. Do not feed to dairy animals or to animals being furnished for slaughter. Keep unprotected persons and beneficial animals out of operating areas. Do not use where there may be danger of drift. Vacated areas should not be re-entered until drifting insecticide and volatile residues have dissipated.

Do not reuse empty container. Destroy it by burying with waste or burning it far away from smokestacks.

Not for use or storage in or around the home.

Birds feeding on treated areas may be killed. Do not contaminate water by use of equipment or disposal of wastes.



**HAVILAND** AGRICULTURAL  
45 STERLING, N.W. BRAND RAPID, M.C. 49114