

# MALATHION COPPER SULFUR DUST NO. 5720

**ACTIVE INGREDIENTS:**

[595-242]

|  |               |
|--|---------------|
| Malathion (O,O-Dimethyl Dithiophosphate of Diethylmercaptosuccinate) ..... | 5.0%          |
| Metallic Copper (from 2-5 micron Tribasic Copper Sulfate) .....            | 7.0%          |
| Sulfur .....   | 20.0%         |
| <b>INERT INGREDIENTS:</b> .....  | <u>68.0%</u>  |
| <b>Total</b>   | <b>100.0%</b> |

Recommended for control of various insects and fungus diseases on Carrots, Celery, Potatoes, Tomatoes, Eggplant and Peppers as suggested below.

**DIRECTIONS:**

The following directions are furnished by us and are believed to be reliable but in no way guaranteed. This dust should be applied at the rates suggested below depending on the size and spacing of the plants and the method of application. **Caution:** Sulfur present in this material may burn foliage when temperatures are extremely high.

**CARROTS** – To Control Aphids and Six Spotted Leaf Hopper, apply dust at the rate of 30 to 40 lbs. per acre. Treat fields and margins when plants are coming up and repeat treatment at 7 day intervals as needed. This treatment will also control Alternaria Leaf Blight. Do not use treated tops for feed, nor apply within 7 days of harvest.

**CELERY** – To control Aphids and Spider Mites and Early and Late Blights when they appear in the field, apply dust at the rate of 25 to 30 lbs. per acre to both the margins and field. Use only very early in the growing season before celery begins to bunch. Repeat application weekly or after a rain as needed. Do not apply within 7 days of harvest.

**POTATOES** – To control Aphids, Leafhoppers, False Chinch

Bug and Mealv Bugs, apply dust at the rate of 30 lbs. per acre when insects first appear. Repeat every 7 to 100 days as needed.

**TOMATOES, EGGPLANT, AND PEPPER** – To control Aphids, Spidermites and Pepper Maggots and Early and Late Blight apply at the rate of 30 to 40 lbs. per acre on tomatoes and eggplant and at the rate of 25 to 30 lbs. per acre on peppers. First treatment should be applied about 30 days after first cluster bloom and sooner if insects appear or blight threatens. Repeat every 7 to 10 days or after a rain if needed. Do not apply within 5 days of harvest on tomatoes and within 3 days of harvest on eggplant or peppers.

**CAUTION: Keep Out of Reach of Children**

Avoid breathing dust. Do not contaminate feed or foodstuffs. Avoid contact with skin, eyes and clothing. In case of contact with skin or eyes, flush with plenty of water; for eyes, get medical attention.

Do not reuse empty container. Destroy it by burying with waste or burning. Stay away from smoke or fumes.

Toxic to fish. Do not contaminate any body of water nor apply when conditions favor drift from the crop area treated. Do not contaminate water by cleaning of equipment, or disposal of wastes.

**NON-WARRANTY**

Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

USDA Reg No. 595-242

A 595-242  
9-16-71 with comments

Net Wt. 50 Lbs.



1345 STERLING AVE. GRAND RAPIDS, MICHIGAN 49502

ACCEPTED WITH COMMENTS