

PARATHION MANEB DUST NO. 17

ACTIVE INGREDIENTS:

Parathion (O,O-Diethyl O-p-nitrophenyl thiophosphate) 1.0%
Maneb (Manganese ethylene bisdithiocarbamate) 7.0%

INERT INGREDIENTS: 92.0%

Total 100.0%



**POISON
ANTIDOTE**



INTERNAL: Give a tablespoon of mustard in a glass of warm water, repeat until vomit fluid is clear. Have victim lie down and remain quiet after the physical vomiting has been carried out. Call a physician immediately. Atropine Sulfate on advice of physician. 2-PAM is also antidotal and may be administered in conjunction with atropine.

EXTERNAL: If skin is contaminated, wash thoroughly with soap and water.

Recommended for control of various insects and fungus diseases on Vegetable Crops as listed below.

DIRECTIONS

As recommendations for the use of Parathion products vary in different agricultural areas, it is suggested that you consult your State Agricultural Experiment Station before using this product. If the crop to be dusted is for canning purposes, consult your cannery fieldman before dusting. The following directions are furnished by us and are believed to be of value. This dust should be applied at the rates suggested below depending on the size and spacing of the plants and the method of application.

CARROTS — To control Aphids, Six Spotted Leafhopper, Alternaria (Leaf Blight) and Cercospora Blight apply 20 to 30 lbs. per acre. Begin application when plants are coming up and repeat weekly as long as insects or blight threatens. In dusting for leafhopper treat both field and field margins. Do not use on carrots within 15 days of harvest. Do not use treated carrot tops for food or feed.

CELERY — To control Aphids, Six Spotted Leafhopper, Tarnish Plant Bug, Early and Late Blight use 30 to 40 lbs. per acre weekly. Begin dusting as soon as insects appear or blight threatens. Do not use after celery begins to bunch or is half grown. Remove excess residues by stripping, trimming, and washing.

CUCUMBER MELON and SQUASH — To control Cucumber Beetle, Aphids, Stink Bug, Melonworm Pickleworm, Anthracnose, Cercospora leaf spot and Downy Mildew apply 30 to 40 lbs. per acre when insects appear or disease threatens. Do not

use on cucumbers before vining, within 15 days of harvest on cucumbers and squash and within 7 days of harvest on melons.

ONIONS — (dry onions only) To control Onion Maggot, Downy Mildew, Blast (Botrytis leaf blight) and Purple Blotch apply 30 to 40 lbs. per acre. For maggot control begin dusting 14 days after planting and repeat at 3 to 5 day intervals as necessary. For fungus control use at weekly intervals when disease is reported in the area. Do not apply within 15 days of harvest.

POTATOES — To control Grasshoppers, Early and Late Blight apply 30 to 50 lbs. per acre when insects appear in large numbers or when blight threatens. Apply every 7 - 10 days throughout the season. Do not apply within 5 days of harvest.

NOTE: For disease control closer to harvest than the given limitations a suitable fungicide should be used alone.

DANGER: Keep Out of Reach of Children

Poisonous if swallowed or inhaled or absorbed through skin or eyes. Rapidly absorbed through skin! Do not get in eyes or on skin or clothing. In case of contact, wash immediately with soap and water. Wear natural rubber gloves, protective clothing and goggles. Wear a mask or respirator of a type passed by U.S. Bureau of Mines for Parathion protection. Wash arms, hands, and face thoroughly after handling and before eating or smoking. Wash all contaminated clothing with soap and hot water before re-use. Keep away from children, animals, feed and foodstuffs. Do not feed to dairy animals or to animals being furnished for slaughter. Keep unprotected persons and beneficial animals out of operating area or vicinity where there may be danger of drift. Vacated areas should not be re entered until drifting insecticide and volatile residues have dissipated. Not for use or storage in or around the home.

Birds feeding on treated areas may be killed. Do not contaminate water by cleaning of equipment, or disposal of wastes. Toxic to fish and wildlife. Do not contaminate any body of water nor apply when weather conditions favor drift from crop area treated. Do not apply to any area not specified on the label.

Do not reuse empty container. Destroy it by burying with wastes or burning. Stay away from smoke or fumes.

NON-WARRANTY

Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

USDA Reg. No. 595-245

9-28-71
595-245

Net Wt. 50 Lbs.