

HAVILAND 3-S DUST

(Contains 3% Sevin*)

ACTIVE INGREDIENTS:

Carbaryl (1-Naphthyl N-methylcarbamate)	3.0%
INERT INGREDIENTS:	97.0%
TOTAL	100.0%

*Sevin is the registered Trade Mark of Union Carbide Corp. for the active ingredient 1-naphthyl N-methylcarbamate.

DIRECTIONS

For specific application consistent with label cautions and directions, it is recommended that you consult your State Agricultural Experiment Station.

BLUEBERRIES – To control Fruitworms, and Fruit Fly apply Haviland 3-S Dust at the rate of 50-70 lbs. per acre. Make first application immediately after bloom or as soon as insects become active and repeat at 10-day intervals until fruit is harvested.

STRAWBERRIES – To control Spittlebug and Leafrollers apply this dust at the rate of 40-60 lbs. per acre for full coverage in regular cover applications. Do not apply within 1 day of harvest.

GRAPES – To control Grape Leaf Folder, Leafhopper and Berry Moth apply at the rate of 40-60 lbs. per acre. Make first application just before first brood leaf folder larvae emerge and repeat at weekly intervals as necessary.

TOMATOES, EGGPLANT and PEPPERS – To control European Corn Borer, Fall Armyworm, Lace Bug, Tomato Fruitworm, Tomato Hornworm, Sinkbugs, Tarnished Plant Bug apply 40-60 lbs. of dust per acre. Make first application when insects or their damage first appear and repeat at 7-10 day intervals as needed

CABBAGE, CAULIFLOWER, BROCCOLI, BRUSSELS SPROUTS and KOHLRABI -- To control Armyworms, Flea Beetle, Harlequin Bug and Corn Earworm apply this dust at the rate of 40-60 lbs. per acre when insects or their damage appear and repeat at 5 to 7 day intervals as necessary. Do not make application within 3 days of harvest.

CAUTION: Keep Out of Reach of Children

Harmful if swallowed or inhaled. Avoid prolonged or repeated contact with skin. Wash thoroughly after using. Avoid breathing dust. Avoid contamination of feed and foodstuffs. Use only on recommended crops and at recommended rates. Avoid storage near feed and foodstuffs.

Do not reuse empty container. Destroy it by burying with waste or burning. Stay away from smoke or fumes.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

Do not contaminate any body of water nor apply when conditions favor drift from the crop area treated.

NON-WARRANTY

Seller makes no warranty, expressed or implied, concerning the use of this product, other than indicated on the label. Buyer assumes all risk of use and/or handling, when such use and/or handling is contrary to label instructions.

USDA Reg. No. 595-247

ACCEPTED
10-17-71
595-247

Net Wt. 50 Lbs.