THIODAN® COPPER DUST

(3 % - 7 %)

ACTIVE INGREDIENTS	
Endosulfan (Hexachlorohexahydromethano 2,4,3 benzodioxathiepin oxide)	3.0%
Metallic Copper (from 2-5 micron Tribasic Copper Sulfate)	7.0%
INERT INGREDIENTS	90.0%
	100.0%

*Thiodan is a registered trademark of Farbwerke Hoechst A.G. for the active ingredient commonly known as endosulfan

DIRECTIONS

The following directions are furnished by us and are believed to be reliable. Make first application when insects appear or disease threatens and repeat weekly if necessary. State agricultural authorities should be consulted for specific timing and dosage rates in each local area.

BEANS To control Mexican Bean Beetle, Downy Mildew, Anthracnose and Angular Leaf Spot, apply 25 to 30 lbs. per acre. Do not apply more than 3 times per season. Do not apply within 3 days of harvest. Do not feed treated threshings to livestock or allow livestock to graze on treated fields. Do not use fume boards that are to be removed from the field for bees.

BROCCOLI, CABBAGE, and CAULIFLOWER To control Cabbage Looper, Imported Cabbage Worm, Diamondback Moth Larvae, Downy Mildew and Leaf Spot apply 25 lbs. per acre. Do not apply within 7 days after edible parts begin to form and do not apply to broccoli or cabbage within 7 days of harvest.

CUCUMBERS, MELONS and SQUASH To control Aphids, Armyworms, Downy Mildew and Angular Leaf Spot apply 25 to 30 lbs. per acre. Apply as needed up to harvest.

HEAD LETTUCE To control Aphids, Cabbage Looper, Imported Cabbage Worm, Diamondback Moth Larvae, and Downy Mildew. Apply 33 to 35 lbs. per acre. Do not make more than 3 applications after thinning. Do not apply within 14 days of harvest. Remove upper leaves at harvest. Do not feed crop refuse to livestock.

PEPPERS and TOMATOES To control Aphids, Colorado Potato Beetle, Flea Beetles, Septoria Leaf Spot and Early and Late Blight, apply 15 to 20 lbs. per acre. Do not apply within 1 day of harvest. When applying in greenhouse wear a mask or respirator passed by U.S. Bureau of Mines for Thiodan protection.

POTATOES To control Aphids, Colorado Potato Beetle, Flea Beetles, Leaf Hoppers, and Early and Late Blight, apply 30 to 35 lbs. per acre. Apply as needed up to harvest. Do not plant root crops other than carrots, potatoes, and sweet potatoes as follow-up crops.

WARNING: Keep Out of Reach of Children

Hazardous if Swallowed, Inhaled or Absorbed Through Skin. Do not breathe dust. Do not get in eyes, on skin, or on clothing. In case of contact, immediately remove contaminated clothing and flush skin or eyes with plenty of water. For eyes get medical attention. Wash thoroughly with soap and water after handling and before eating or smoking. Wear clean clothing. During operations involving prolonged exposure, spray mixing and loading operations, wear mask or respirator of a type passed by the U.S. Bureau of Mines for endosulfan protection.

Birds feeding on treated areas may be killed. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not apply where runoff is likely to contaminate any body of water. Do not apply when weather conditions favor drift from treated areas. Do not apply in any manner not specified on this label.

Do not reuse empty containers. Triple rinse by filling container 1/4 full with water and shake for 5 seconds. Pour rinsate into application equipment or store rinsate for later use.

NON-WARRANTY

Seller makes no warranty, express or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and handling of this material when such use and handling is contrary to label directions.

USDA Reg. No. 595-248

9-21-71
595-248

Net Wt. 50 Lbs